

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



B11

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 7/10, A61K 37/43	A1	(11) International Publication Number: WO 94/01460 (43) International Publication Date: 20 January 1994 (20.01.94)
(21) International Application Number: PCT/US93/06465 (22) International Filing Date: 13 July 1993 (13.07.93) (30) Priority data: 07/915,247 14 July 1992 (14.07.92) US (71) Applicant: SYNTEX (U.S.A.) INC. [US/US]; 3401 Hillview Avenue A2-200, Palo Alto, CA 94304 (US). (72) Inventors: KRSTENANSKY, John, L. ; 3455 Rambo Drive, Palo Alto, CA 94306 (US). NESTOR, John, J., Jr. ; 20937 Fairwoods Court, Cupertino, CA 95014 (US). HO, Teresa, L. ; 495 South Clark Avenue, Los Altos, CA 94024 (US). VICKERY, Brian, H. ; 111 Woofleaf Way, Mountain View, CA 94040 (US). BACH, Chinh, T. ; 1613 Adolfo Drive, San Jose, CA 95131 (US).		(74) Agent: CHOW, Y., Ping; Syntex (U.S.A.) Inc., 3401 Hillview Avenue A2-200, Palo Alto, CA 94304 (US). (81) Designated States: AU, CA, FI, HU, JP, KR, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ANALOGS OF PTH AND PTHRP, THEIR SYNTHESIS AND USE FOR THE TREATMENT OF OSTEOPOROSIS (57) Abstract Synthetic polypeptide analogs of parathyroid hormone PTH, parathyroid hormone related peptide PTHrp, and of the physiologically active truncated homologs and analogs of PTH and PTHrp, in which amino acid residues (22-31) form an amphipathic α -helix, said residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa) ordered in the sequence: Haa(Laa Laa Haa Haa) ₂ Laa and their pharmaceutically acceptable salts are useful for the prophylaxis and treatment of osteoporosis in mammals. Processes for the production of the polypeptides via solid phase and recombinant methods are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

5 ANALOGS OF PTH AND PTHRP , THEIR SYNTHESIS AND USE FOR THE TREATMENT OF
OSTEOPOROSIS.

BACKGROUND OF THE INVENTION

10

a) Field of the Invention

15 This invention relates to novel analogs of parathyroid hormone and parathyroid hormone related peptide, their synthesis by solid phase and recombinant techniques, and their use for increasing bone mass in mammalian subjects.

b) Description of Related Art

20

Osteoporosis is the most common form of metabolic bone disease and may be considered the symptomatic, fracture stage of bone loss (osteopenia). Although osteoporosis may occur secondary to a number of underlying diseases, 90% of all cases appear to be idiopathic. Postmenopausal women are particularly at risk for idiopathic osteoporosis (postmeno-pausal or Type I osteoporosis). Another high risk group for

25

idiopathic osteoporosis is the elderly of either sex (senile or Type II osteoporosis). Osteoporosis has also been related to corticosteroid use, immobilization or extended bed rest, alcoholism, diabetes, gonadotoxic chemotherapy, hyperprolactinemia, anorexia nervosa, primary and secondary

30

amenorrhea, and oophorectomy. In the various forms of osteoporosis, bone fractures, which are the result of bone loss that has reached the point of mechanical failure, frequently occur. Postmenopausal osteoporosis is characterized by fractures of the wrist and spine, while femoral neck fractures seem to be the dominant feature of senile osteoporosis.

35

The mechanism by which bone is lost in osteoporotics is believed to involve an imbalance in the process by which the skeleton renews itself.

40

This process has been termed bone remodeling. It occurs in a series of discrete pockets of activity. These pockets appear spontaneously within the bone matrix on a given bone surface as a site of bone resorption. Osteoclasts (bone dissolving or resorbing cells) are responsible for the resorption of a portion of bone of generally constant dimension. This resorption process is followed by the appearance of osteoblasts (bone forming cells) which then refill with new bone the cavity left by the osteoclasts.

45

In a healthy adult subject, the rate at which osteoclasts and osteoblasts are formed is such that bone formation and bone resorption are in balance. However, in osteoporotics an imbalance in the bone remodeling process develops which results in bone being lost at a rate faster than it

is being made. Although this imbalance occurs to some extent in most individuals as they age, it is much more severe and occurs at a younger age in postmenopausal osteoporotics or following oophorectomy.

5 There have been many attempts to treat osteoporosis with the goal of either slowing further bone loss or, more desirably, producing a net gain in bone mass. Certain agents, such as estrogen and the bisphosphonates, appear to slow further bone loss in osteoporotics. Agents which slow bone loss, because of the different durations of bone resorption and formation, may appear to increase bone mass (on the order of 3 to 7%). However, this
10 apparent increase is limited in time, not progressive, and is due to a decrease in "remodeling space." In addition, because of the close coupling between resorption and formation, treatments which impede bone resorption also ultimately impede bone formation.

15 It has been suggested that treatment with parathyroid hormone (PTH) would lead to both increased bone turnover and a positive calcium balance. However, human clinical trials have shown that any increase in trabecular bone is offset by a decrease in cortical bone, so that there is no net increase in total bone.

20 Hefti, et al. in *Clinical Science* 62, 389-396 (1982) have reported that daily subcutaneous doses of either bPTH(1-84) or hPTH(1-34) increased whole body calcium and ash weight of individual bones in both normal and osteoporotic adult female rats.

25 Liu, et al. in *J. Bone Miner. Res.* 6:10, 1071-1080 (1991) have noted that ovariectomy of adult female rats induced a 47% loss in the percentage of trabecular bone in the proximal tibial metaphysis, accompanied by a significant increase in the number of osteoblasts and trabecular osteoclasts. Daily subcutaneous injections of hPTH(1-34) completely reversed the loss of trabecular bone and resulted in amounts of trabecular bone exceeding that of sham operated controls. The number of osteoblasts
30 increased and the number of osteoclasts decreased.

Hock et al. in *J. Bone Min. Res.* 7:1, 65-71 (1992) have reported that daily subcutaneous injections of hPTH(1-34) to healthy adult male rats for 12 days increased trabecular and cortical bone calcium and dry weight. Total bone mass, trabecular bone volume, trabecular thickness and number, and osteoblastic surfaces were increased.
35

The mammalian parathyroid hormones, e.g. human (hPTH), bovine (bPTH), and porcine (pPTH), are single polypeptide chains of 84 amino acid residues, with molecular weights of approximately 9500. Biological activity is associated with the N-terminal portion, with residues (1-34) apparently the minimum required.
40

The N-terminal segment of human PTH differs from the N-terminal segment of the bovine and porcine hormones by only three and two amino acid residues, respectively:

-3-

hPTH(1-34):

Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu
 1 5 10 15

Asn Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp
 20 25 30

Val His Asn Phe (SEQ ID NO:1);

bPTH(1-34):

Ala Val Ser Glu Ile Gln Phe Met His Asn Leu Gly Lys His Leu
 1 5 10 15

Ser Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp
 20 25 30

Val His Asn Phe (SEQ ID NO:2);

pPTH(1-34):

Ser Val Ser Glu Ile Gln Leu Met His Asn Leu Gly Lys His Leu
 1 5 10 15

Ser Ser Leu Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp
 20 25 30

Val His Asn Phe (SEQ ID NO:3).

The primary function of PTH is to elicit the adaptive changes that serve to maintain a constant concentration of Ca^{2+} in the extracellular fluid. PTH acts on the kidneys to increase tubular reabsorption of Ca^{2+} from the urine, as well as stimulating the conversion of calcifediol to calcitriol, which is responsible for absorption of Ca^{2+} from the intestines. One prominent effect is to promote the mobilization of Ca^{2+} from bone. PTH acts on bone to increase the rate of resorption of Ca^{2+} and phosphate. PTH stimulates the rate of bone resorption by osteoclasts, increases the rate of differentiation of mesenchymal cells to osteoclasts, and prolongs the half-life of these latter cells. With prolonged action of PTH the number of bone forming osteoblasts is also increased; thus, the rate of bone turnover and remodeling is enhanced. However, individual osteoblasts appear to be less active than normal.

Rosenblatt, et al. in U.S. Patent Nos. 4,423,037, 4,968,669 and 5,001,223 have disclosed PTH antagonists obtained by the deletion of the N-terminal (1-6) amino acids and the selective replacement of Phe⁷, Met^{8,18}, and Gly¹². Tyr³⁴-NH₂ reportedly increased the activity and stability of these compounds.

Parathyroid hormone-related peptide (PTHrp), a 140+ amino acid protein, and fragments thereof, reproduce the major biological actions of PTH. PTHrp is elaborated by a number of human and animal tumors and other tissues and may play a role in hypercalcemia of malignancy. The sequence of hPTHrp (1-34) is as follows:

-4-

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 5 Gln Asp Leu Arg Arg Arg Phe Phe Leu His His Leu Ile Ala Glu
 20 25 30
 Ile His Thr Ala (SEQ ID NO:4).

10 The sequence homology between hPTH and hPTHrp is largely limited to the 13 N-terminal residues, 8 of which are identical; only 1 of 10 amino acids in the (25-34) receptor binding region of hPTH is conserved in hPTHrp. Conformational similarity may underlie the common activity. Cohen, et al. in *J. Biol. Chem.* 266:3, 1997-2004 (1991) have suggested that
 15 much of the sequence of PTH(1-34) and PTHrp(1-34), in particular regions (5-18) and (21-34), assumes an α -helical configuration, while noting that there is some question whether this configuration prevails for the carboxyl terminal end under physiological conditions. Such a secondary structure may be important for lipid interaction, receptor interaction, and/or structural stabilization.

20 We have synthesized analogs of PTH and of PTHrp with the objective of developing improved therapeutic agents for the restoration of bone mass in mammalian subjects, including those afflicted with osteoporosis.

25 SUMMARY OF THE INVENTION

This invention provides synthetic polypeptide analogs of parathyroid hormone (PTH), parathyroid hormone related peptide (PTHrp), and of the physiologically active truncated homologs and analogs of PTH and PTHrp, and salts thereof, in which amino acid residues (22-31) form an amphipathic
 30 α -helix, said residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa) ordered in the sequence:

Haa (Laa Laa Haa Haa)₂ Laa.

35 When specific illustrative embodiments of this sequence type are inserted into PTH, PTHrp, and the physiologically active truncated analogs and homologs of PTH and PTHrp, the resulting polypeptides are effective bone remodeling agents.

In one aspect, then, this invention provides analogs of PTH, PTHrp, and the physiologically active truncated analogs and homologs of PTH and PTHrp, or salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, the sequence of said residues (22-31) selected from:

a) Xaa¹ Xaa² Leu Xaa⁴ Xaa⁵ Leu Xaa⁷ Xaa⁸ Xaa⁹ Xaa¹⁰ wherein
 1 5 10

45 Xaa¹ and Xaa⁴ are independently Glu, Glu(OCH₃), His, or Phe; Xaa² is Leu or Phe; Xaa⁵ is Lys or His; Xaa⁷ and Xaa¹⁰ are independently Leu or Ile; Xaa⁸ is Ala, Arg, or Glu; and Xaa⁹ is Lys or Glu (SEQ ID NO:85); preferably

-5-

Glu Leu Leu Glu Lys Leu Leu Xaa Lys Leu wherein
 1 5 10

Xaa is Glu or Arg (SEQ ID NO:26);

5 b) Xaa¹ Xaa² Leu Xaa⁴ Arg Leu Leu Xaa⁸ Arg Leu wherein
 1 5 10

Xaa¹ and Xaa⁴ are independently Glu, Glu(OCH₃), His, or Phe; Xaa² is Leu or Phe; Xaa⁸ is Glu, Lys, or Lys(COCH₂PEGX) and PEGX is a poly-(ethylene glycol methyl ether) radical of molecular weight 100 to 10,000 (SEQ ID NO:86); preferably,

10 Glu Leu Leu Glu Arg Leu Leu Xaa Arg Leu wherein
 1 5 10

Xaa is Glu, Lys, or Lys(COCH₂PEGX) and PEGX is a poly-(ethylene glycol methyl ether) radical of molecular weight 100 to 10,000 (SEQ ID NO:27);

15 c) Ala Leu Ala Glu Ala Leu Ala Glu Ala Leu (SEQ ID NO:28);
 1 5 10

d) Ser Leu Leu Ser Ser Leu Leu Ser Ser Leu (SEQ ID NO:29);
 1 5 10

20 e) Ala Phe Tyr Asp Lys Val Ala Glu Lys Leu (SEQ ID NO:30).
 1 5 10

In another aspect, this invention provides analogs of PTH, PTHrp, and of the physiologically active truncated homologs and analogs of PTH and PTHrp, or salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, the sequence of said residues (22-31) selected from:

25 a) Glu Leu Leu Glu Lys Leu Leu Xaa Lys Leu wherein
 1 5 10

Xaa is Glu or Arg (SEQ ID NO:26);

30 b) Glu Leu Leu Glu Arg Leu Leu Xaa Arg Leu wherein
 1 5 10

Xaa is Glu, Lys, or Lys(COCH₂PEGX) and PEGX is a poly-(ethylene glycol methyl ether) radical of molecular weight 100 to 10,000 (SEQ ID NO:27);

c) Ala Leu Ala Glu Ala Leu Ala Glu Ala Leu (SEQ ID NO:28);
 1 5 10

35 d) Ser Leu Leu Ser Ser Leu Leu Ser Ser Leu (SEQ ID NO:29);
 1 5 10

e) Ala Phe Tyr Asp Lys Val Ala Glu Lys Leu (SEQ ID NO:30).
 1 5 10

40 Also provided are pharmaceutical compositions for the prevention or treatment of conditions characterized by decreases in bone mass comprising an effective bone mass increasing amount of a polypeptide analog of parathyroid hormone (PTH), parathyroid hormone related peptide (PTHrp), and of the physiologically active truncated homologs and analogs of PTH and PTHrp, and salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, said residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa) ordered in the sequence:

Haa (Laa Laa Haa Haa), Laa.

in admixture with a pharmaceutically acceptable carrier. Also provided are processes for preparing pharmaceutical compositions which comprise mixing the above described compounds with a pharmaceutically acceptable carrier.

5 Further, this invention provides methods for treating mammalian conditions characterized by decreases in bone mass which methods comprise administering to a mammal in need thereof an effective bone mass increasing amount of a polypeptide analog of PTH, PTHrp, or of a physiologically active truncated homolog or analog of PTH or PTHrp, or a salt thereof, in which amino acid residues (22-31) form an amphipathic α -helix, said
10 residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa) ordered in the sequence:

Haa (Laa Laa Haa Haa)₂ Laa.

15 More specifically, this invention provides methods for treating mammalian conditions characterized by decreases in bone mass which methods comprise administering to a mammal in need thereof an effective bone mass increasing amount of a polypeptide analog of PTH, PTHrp, or of a physiologically active truncated homolog or analog of PTH or PTHrp, or salt thereof, in which amino acid residues (22-31) form an amphipathic α -helix,
20 the sequence of said residues (22-31) selected from: (SEQ ID NOS: 26, 27, 28, 29, and 30).

This invention also includes processes for the solid phase synthesis of polypeptide analogs of PTH, PTHrp, and of the physiologically active truncated homologs and analogs of PTH and PTHrp, and salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, said
25 residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa) ordered in the sequence:

30 Haa (Laa Laa Haa Haa)₂ Laa,

which processes comprise sequentially coupling protected amino acids on a suitable resin support, removing the side chain and N^o-protecting groups, and cleaving the polypeptide from the resin.

35 This invention also encompasses processes for the solid phase synthesis of polypeptide analogs of PTH, pTHrp, and of the physiologically active truncated homologs and analogs of PTH and PTHrp, and salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, the sequence of said residues (22-31) selected from: (SEQ ID NOS: 26, 27, 28,
40 29, and 30); which processes comprise sequentially coupling protected amino acids on a suitable resin support, removing the side chain and N^o-protecting groups, and cleaving the polypeptide from the resin.

Also included are processes for the recombinant synthesis of polypeptide analogs of PTH, PTHrp, and of the physiologically active

- 7 -

truncated homologs and analogs of PTH and PTHrp, and salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, said residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa) ordered in the sequence:

Haa (Laa Laa Haa Haa)₂ Laa.

The invention also encompasses DNA sequences, vectors, and plasmids for the recombinant synthesis of such polypeptide analogs. Specifically, the invention provides DNA sequences, vectors, and plasmids for the recombinant synthesis of polypeptide analogs of PTH, PTHrp, and of the physiologically active truncated homologs and analogs of PTH and PTHrp, and salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, the sequence of said residues (22-31) selected from: (SEQ ID NOS: 26, 27, 28, 29, and 30).

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 presents the DNA sequence and enzyme restriction sites of a synthetic gene coding for a PTHrp(1-34) analog of this invention (hPTH PCR Amplification).

Figure 2 outlines the preparation of a plasmid incorporating a PTHrp(1-34) analog gene (Trp LE 18 hPTHrp(1-34)1 Construction).

Figure 3 outlines the preparation of a plasmid incorporating two copies of a PTHrp(1-34) analog gene (Trp LE 18 hPTHrp(1-34)2 Construction).

Figure 4 outlines the preparation of a plasmid incorporating four copies of a PTHrp(1-34) analog gene (Trp LE 18 hPTHrp(1-34)4 Construct).

DETAILED DESCRIPTION OF THE INVENTION

Abbreviations and Definitions

The one- and three-letter abbreviations for the various common nucleotide bases and amino acids are as recommended in *Pure Appl. Chem.* 31, 639-645 (1972) and 40, 277-290 (1974) and comply with 37 CFR §1.822 (55 FR 18245, May 1, 1990), and with the PCT rules (WIPO Standard ST.23: *Recommendation for the Presentation of Nucleotide and Amino Acid Sequences in Patent Applications and in Published Patent Documents*). The one- and three-letter abbreviations are as follows:

**SUBSTITUTE SHEET
ISA/EP**

- 8 -

<u>Amino Acid Abbreviations</u>			
	Amino Acid	Three-letter Symbol	One-letter Symbol
5	Alanine	Ala	A
	Arginine	Arg	R
	Asparagine	Asn	N
	Aspartic Acid	Asp	D
	Asn + Asp	Asx	B
10	Cysteine	Cys	C
	Glutamine	Gln	Q
	Glutamic Acid	Glu	E
	Gln + Glu	Glx	Z
	Glycine	Gly	G
15	Histidine	His	H
	Isoleucine	Ile	I
	Leucine	Leu	L
	Lysine	Lys	K
	Methionine	Met	M
20	Phenylalanine	Phe	F
	Proline	Pro	P
	Serine	Ser	S
	Threonine	Thr	T
	Tryptophan	Trp	W
25	Tyrosine	Tyr	Y
	Valine	Val	V
	Other amino acid	Xaa	X

The abbreviations represent L-amino acids unless otherwise designated as D- or D,L-. Certain amino acids, both natural and non-natural, are achiral, e.g. glycine. All peptide sequences are presented with the N-terminal amino acid on the left and the C-terminal amino acid on the right.

Further abbreviations for other amino acids and compounds used herein are:

hSer	homoserine
hSerlac	homoserine lactone
Nle	norleucine
PEG2	radical of diethylene glycol methyl ether, a.k.a. methoxydi(ethyleneoxy), $\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_2^-$, (MW = 119)
PEG5000	radical of poly(ethylene glycol methyl ether), a.k.a. methoxypoly(ethyleneoxy), $\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_{110}^-$, (avg. MW = 5000)
PEGX	radical of poly(ethylene glycol methyl ether), $\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_n^-$, $n = 2 - 225$, (avg. MW = 100 to 10,000)

"Hydrophilic amino acid (Haa)" refers to an amino acid having at least one hydrophilic functional group in addition to those required for peptide bond formation, such as arginine, asparagine, aspartic acid, glutamic acid, glutamine, histidine, lysine, serine, threonine, and their homologs.

"Lipophilic amino acid (Laa)" refers to an uncharged, aliphatic or aromatic amino acid, such as isoleucine, leucine, methionine, phenylalanine, tryptophan, tyrosine, valine, and their homologs.

For the purposes of this invention, alanine is classified as "amphiphilic" i.e., capable of acting as either hydrophilic or lipophilic.

"Physiologically active truncated homolog or analog of PTH or PTHrp" refers to a polypeptide having a sequence comprising less than the full complement of amino acids found in PTH or PTHrp which, however, elicits a similar physiological response. The truncated PTH or PTHrp need not be fully homologous with PTH or PTHrp to elicit a similar physiological response. PTH(1-34) and PTHrp(1-34) are preferred, but not exclusive, representatives of this group.

"Amphipathic α -helix" refers to the secondary structure exhibited by certain polypeptides in which the amino acids assume an α -helical configuration having opposing polar and nonpolar faces oriented along the long axis of the helix. The possibility of α -helical structure in the polypeptide of interest may be explored to some extent by the construction of a "Schiffer-Edmundson wheel" (M. Schiffer and A. B. Edmundson, *Biophys. J.* 7, 121 (1967)), of the appropriate pitch and noting the segregation of the hydrophilic and lipophilic residues on opposite faces of the cylinder circumscribing the helix. Alternatively, empirical evidence, such as circular dichroism or x-ray diffraction data, may be available indicating the presence of an α -helical region in a given polypeptide. An ideal α -helix has 3.6 amino acid residues per turn with adjacent side chains separated by 100° of arc. Eisenberg et al. in *Nature* 299:371-374 (1982) and *Proc. Nat. Acad. Sci. USA* 81:140-144 (1984) have combined a hydrophobicity scale with the helical wheel to quantify the concept of amphipathic helices. The mean hydrophobic moment is defined as the vector sum of the hydrophobicities of the component amino acids making up the helix. The following hydrophobicities for the amino acids are those reported by Eisenberg (1984) as the "consensus" scale:

Ile 0.73; Phe 0.61; Val 0.54; Leu 0.53; Trp 0.37;
Met 0.26 Ala 0.25; Gly 0.16; Cys 0.04; Tyr 0.02;
Pro -0.07; Thr -0.18; Ser -0.26; His -0.40; Glu -0.62;
Asn -0.64; Gln -0.69; Asp -0.72; Lys -1.10; Arg -1.76.

The hydrophobic moment, μ_H , for an ideal α -helix having 3.6 residues per turn (or a 100° arc ($= 360^\circ/3.6$) between side chains), may be calculated from:

$$\mu_H = [(\sum H_N \sin \delta (N-1))^2 + (\sum H_N \cos \delta (N-1))^2]^{1/2},$$

where H_N is the hydrophobicity value of the N^{th} amino acid and the sums are taken over the N amino acids in the sequence with periodicity $\delta=100^\circ$. The hydrophobic moment may be expressed as the mean hydrophobic moment per residue by dividing μ_H by N to obtain $\langle\mu_H\rangle$. A value of $\langle\mu_H\rangle$ at $100^\circ \pm 20^\circ$ of about 0.20 or greater is suggestive of amphipathic helix formation. The $\langle\mu_H\rangle$ values at 100° for hPTHrp (22-31) and hPTH (22-31) are 0.19 and 0.37, respectively.

Cornett, et al., in *J. Mol. Biol.*, 195:659-685 (1987) have further extended the study of amphipathic α -helices by introducing the "amphipathic index" as a predictor of amphipathicity. They concluded that approximately half of all known α -helices are amphipathic, and that the dominant frequency is 97.5° rather than 100° , with the number of residues per turn being closer to 3.7 than 3.6. While such refinements are scientifically interesting, the basic approach of Eisenberg, et al. is sufficient to classify a given sequence as amphipathic, particularly when one is designing a sequence *ab initio* to form an amphipathic α -helix.

A substitute amphipathic α -helical amino acid sequence may lack homology with the sequence of a given segment of a naturally occurring polypeptide but elicits a similar secondary structure, i. e. an α -helix having opposing polar and nonpolar faces, in the physiological environment. Replacement of the naturally occurring amino acid sequence with an alternative sequence may beneficially affect the physiological activity, stability, or other properties of the altered parent polypeptide. Guidance as to the design and selection of such sequences is provided in J. L. Krstenansky, et al., *FEBS Letters* 242:2, 409-413 (1989), and J. P. Segrest, et al. *Proteins: Structure, Function, and Genetics* 8:103-117 (1990) among others.

The ten amino acid amphipathic α -helix of this invention has the formula:

Haa (Laa Laa Haa Haa), Laa

wherein the Haa's are selected from the group of hydrophilic amino acids and the Laa's are selected from the group of lipophilic amino acids, as defined above. Assuming an idealized α -helix, residues 1, 4, 5, 8, and 9 are distributed along one face (A) of the helix within about a 140° arc of each other, while residues 2, 3, 6, 7, and 10 occupy an opposing 140° arc on the other face (B) of the helix. Preferably, all the residues on one face are of the same polarity while all those on the other face are of the opposite polarity, i.e., if face A is all hydrophilic, face B is all lipophilic and vice versa. The skilled artisan will recognize that while the helices of this invention are described by

Haa(Laa Laa Haa Haa), Laa,

the reverse sequence,

-11-

Laa (Haa Haa Laa Laa)₂ Haa

will also meet the residue distribution criteria and is an equivalent descriptor of the helices of this invention.

5 Alanine may be substituted for either hydrophilic or lipophilic amino acids, since Ala can reside readily on either face of an amphipathic α -helix, although Ala₁₀ does not form an amphipathic α -helix. Generally, proline, cysteine, and tyrosine are not used; however, their presence and other random errors in the sequence may be tolerated, e.g. a hydrophilic
10 residue on the lipophilic face, as long as the remaining amino acids in the segment substantially conform to the hydrophilic face - lipophilic face division. A convenient method for determining if a sequence is sufficiently amphipathic to be a sequence of this invention is to calculate the mean hydrophobic moment, as defined above. If the peak mean moment per
15 residue at $100^\circ \pm 20^\circ$ exceeds about 0.20, then the sequence will form an amphipathic helix and is a sequence of this invention.

For example, the mean hydrophobic moment per residue at 100° for (SEQ ID NO: 26), Xaa = Glu, is calculated as follows:

	A.A.	H _N	$\delta(N-1)$	$H \sin \delta(N-1)$	$H \cos \delta(N-1)$
20	E	-.62	0	0	-.62
	L	.53	100	.52	-.17
	L	.53	200	-.18	-.50
	E	-.62	300	.34	-.31
	K	-1.1	400	-.70	-.85
25	L	.53	500	.34	-.41
	L	.53	600	-.46	-.27
	E	-.62	700	.21	-.58
	K	-1.1	800	-1.08	-.19
	L	.53	900	0	-.53
30				$\Sigma=0.81$	$\Sigma=-4.43$

$$\mu_H = [(0.81)^2 + (-4.43)^2]^{1/2} = 4.50$$

35

$$< \mu_H > = 4.50/10 = 0.45$$

For this sequence, the mean peak hydrophobic moment occurs at 92° and has a value of 0.48.

40 In applying this concept to parathyroid hormone and parathyroid hormone related peptide, it was hypothesized that either or both regions (7-16) and (22-31) may exhibit α -helical secondary structure and could be replaced with a non-homologous sequence having similar structural tendencies, without loss of biological activity or induction of immunoreaction.

In one aspect, this invention provides analogs of PTH, PTHrp, and the physiologically active truncated analogs and homologs of PTH and PTHrp, or salts thereof, in which amino acid residues (22-31) form an amphipathic α -helix, the sequence of said residues (22-31) selected from:

Xaa¹ and Xaa⁴ are independently Glu, Glu(OCH₃), His, or Phe; Xaa² is Leu or Phe; Xaa³ is Lys or His; Xaa⁷ and Xaa¹⁰ are independently Leu or Ile; Xaa⁸ is Ala, Arg, or Glu; and Xaa⁹ is Lys or Glu (SEQ ID NO: 85); preferably
Glu Leu Leu Glu Lys Leu Leu Xaa Lys Leu wherein
_{1 5 10}

Xaa¹ and Xaa⁴ are independently Glu, Glu(OCH₃), His, or Phe; Xaa² is Leu or Phe; Xaa⁸ is Glu, Lys, or Lys(COCH₂PEGX) and PEGX is a poly-(ethylene glycol methyl ether) radical of molecular weight 100 to 10,000 (SEQ ID NO:86); preferably, Glu₁ Leu₁ Leu₅ Glu Arg Leu Leu Xaa Arg Leu wherein

e) Ala Phe Tyr Asp Lys Val Ala Glu Lys Leu (SEQ ID NO:30).
1 5 10

Leu Xaa¹⁹ Xaa²⁰ Xaa²¹ Xaa²²⁻³¹ Xaa³² Xaa³³ Xaa³⁴ Xaa³⁵ Xaa³⁶ Xaa³⁷ Xaa³⁸ Term, wherein:

Xaa¹³ is Lys, Arg, Tyr, Cys, Leu, Cys(CH₂CONH(CH₂)₂NH(biotinyl)), Lys(7-dimethylamino-2-oxo-2H-1-benzopyran-4-acetyl), or Lys(dihydrocinnamoyl);

-13-

Xaa²⁰ is Arg or Leu;

Xaa¹⁹ and Xaa²¹ are independently Lys, Ala, or Arg;

Xaa²²⁻³¹ is selected from (SEQ ID NOS:26, 27, 28, 29, or 30); Xaa³² is His, Pro, or Lys;

5 Xaa³³ is absent, or is Pro, Thr, Glu, or Ala;

Xaa³⁴ is absent, or is Pro, Arg, Met, Ala, hSer, hSer lactone, Tyr, Leu, or 1,4-diaminobutyryl lactam;

Xaa³⁵ is absent or is Pro, Glu, Ser, Ala, or Gly;

Xaa³⁶ is absent or is Ala, Arg, or Ile;

10 Xaa³⁷ is absent or is Arg, Trp, or 3-(-2-naphthyl)-L-alanine;

Xaa³⁸ is absent or is Ala or hSer or Xaa³⁸⁻⁴² is Thr Arg Ser Ala Trp; and Term is OR or NR₂ where each R is independently H, (C₁-C₄)alkyl or phenyl(C₁-C₄)alkyl; and the pharmaceutically acceptable salts thereof.

15 In yet another aspect this invention includes polypeptide analogs of the physiologically active truncated homolog hPTHrp(1-34), as shown in Formula (I):

Ala Val Ser Glu Xaa⁵ Gln Leu Leu His Asp Xaa¹¹ Gly Xaa¹³ Ser Ile Gln Asp Leu Xaa¹⁹ Arg Xaa²¹ Xaa²²⁻³¹ Xaa³² Xaa³³ Xaa³⁴ Term, wherein:

Xaa⁵ is His or Ala;

20 Xaa¹¹ and Xaa¹³ are independently Lys, Arg, or Leu;

Xaa¹⁹ and Xaa²¹ are independently Ala or Arg;

Xaa²²⁻³¹ is selected from:

a) Glu Leu Leu Glu Lys Leu Leu Xaa Lys Leu wherein
1 5 10

25 Xaa is Glu or Arg (SEQ ID NO:26);

b) Glu Leu Leu Glu Arg Leu Leu Xaa Arg Leu wherein
1 5 10

30 Xaa is Glu, Lys, or Lys(COCH₂PEGX) and PEGX is a poly(ethylene glycol methyl ether) radical of molecular weight 100 to 10,000 (SEQ ID NO:27);

~~c) Ala Leu Ala Glu Ala Leu Ala Glu Ala Leu (SEQ ID NO:28);~~
1 5 10

40 d) Ser Leu Leu Ser Ser Leu Leu Ser Ser Leu (SEQ ID NO:29);
1 5 10

e) Ala Phe Tyr Asp Lys Val Ala Glu Lys Leu (SEQ ID NO:30);
1 5 10

45 Xaa³² is His or Lys;

Xaa³³ is Thr, Glu, or Ala;

Xaa³⁴ is Ala, hSer, Tyr, or Leu;

50 and Term is Gly Arg Arg, lactone, OH or NR₂, where each R is H or (C₁-C₄) alkyl; and their pharmaceutically acceptable salts. (Formula I)

A more specific aspect of the invention includes

those polypeptides of Formula (I) wherein Xaa²²⁻³¹ is (SEQ ID NO:26), for which $\langle \mu_H \rangle$ at 100° exceeds 0.45. A still more specific aspect of the invention includes those Formula (I) polypeptides wherein Xaa²²⁻³¹ is (SEQ ID NO:26); Xaa¹¹ and Xaa¹³ are both Lys; and Xaa¹⁹ and Xaa²¹ are both Arg.

Representative polypeptides include, but are not limited to:

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Arg Lys
20 25 30

Leu His Thr Ala OH (SEQ ID NO:5);

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Thr Ala OH (SEQ ID NO:6);

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Thr Ala NH₂ (SEQ ID NO:7);

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Thr hSer NH₂ (SEQ ID NO:8);

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Thr hSerlac (SEQ ID NO:9);

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Thr Ala Gly Arg Arg OH (SEQ ID NO:10); and
35

-15-

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 5 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu Lys Glu Leu NH₂ (SEQ ID NO:11).

Another aspect of this invention includes those polypeptides of Formula
 10 (I) wherein Xaa²³¹ is (SEQ ID NO:26); Xaa¹¹ and Xaa¹³ are both Lys; and one
 of Xaa¹⁹ and Xaa²¹ is Arg and the other is Ala. Representative poly-peptides
 of this subgenus include, but are not limited to:

15 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Ala Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 20 25 30
 Leu His Thr Ala NH₂ (SEQ ID NO:12) and
 25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Ala Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 30 Leu His Thr Ala NH₂ (SEQ ID NO:13).

In another aspect this invention includes those polypeptides of
 Formula (I) wherein Xaa²³¹ is (SEQ ID NO:26); one of Xaa¹¹ and Xaa¹³ is Leu
 35 and the other is Lys; and Xaa¹⁹ and Xaa²¹ are both Arg. Representative poly-
 peptides of this subgenus include, but are not limited to:

Ala Val Ser Glu Ala Gln Leu Leu His Asp Leu Gly Lys Ser Ile
 1 5 10 15
 40 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Ala Leu OH (SEQ ID NO:14).

45 In another aspect this invention includes those polypeptides of Formula
 (I) wherein Xaa²³¹ is (SEQ ID NO:27), for which $\langle \mu_H \rangle$ at 100° exceeds 0.50.
 A further aspect of this invention includes those Formula (I) polypeptides
 wherein Xaa²³¹ is (SEQ ID NO:27); Xaa¹¹ and Xaa¹³ are both Lys or both Arg;
 50 and Xaa¹⁹ and Xaa²¹ are both Arg. Representative polypeptides of this
 subgenus include, but are not limited to:

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 55 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30
 Leu His Thr Ala OH (SEQ ID NO:15);

-16-

1 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 5 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30
 Leu His Thr Ala OH (SEQ ID NO:16);
 10 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 15 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Lys Arg
 20 25 30
 Leu His Thr Ala OH (SEQ ID NO:17);
 20 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu
 20 25
 25 Lys(COCH₂PEG2) Arg Leu His Thr Ala OH (SEQ ID NO:18); and
 30
 30 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu
 20 25
 35 Lys(COCH₂PEG5000) Arg Leu His Thr Ala OH (SEQ ID NO:19).
 30

40 In another aspect this invention includes polypeptides of Formula (I)
 wherein Xaa²²⁻³¹ is (SEQ ID NO:28), for which $\langle \mu_H \rangle$ at 100° is about 0.25.
 Representative polypeptides of this subgenus include, but are not limited
 to:

45 1 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Ala Leu Ala Glu Ala Leu Ala Glu Ala
 20 25 30
 50 Leu His Thr Ala NH₂ (SEQ ID NO:20).

55 In another aspect this invention includes polypeptides of Formula (I)
 wherein Xaa²²⁻³¹ is (SEQ ID NO:29), for which $\langle \mu_H \rangle$ at 100° is about 0.28.
 Representative polypeptides of this subgenus include, but are not limited
 to:

60 1 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Ser Leu Leu Ser Ser Leu Leu Ser Ser
 20 25 30
 Leu His Thr Ala NH₂ (SEQ ID NO:21).

-17-

In another aspect this invention includes polypeptides of Formula (I) wherein Xaa²²⁻³¹ is (SEQ ID NO:30), for which $\langle \mu_H \rangle$ at 100° is about 0.29. Representative polypeptides of this subgenus include, but are not limited to:

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
10 Gln Asp Leu Arg Arg Arg Ala Phe Tyr Asp Lys Val Ala Glu Lys
 20 25 30
Leu His Thr Ala NH₂ (SEQ ID NO:22).

15

Still another aspect of this invention includes polypeptide analogs of the physiologically active truncated homolog bPTH(1-34), as shown in Formula (II):

20 Xaa¹ Val Ser Glu Ile Gln Xaa⁷ Xaa⁸ His Asn Leu Gly Lys His Leu Xaa¹⁶ Ser Xaa¹⁸
Xaa¹⁹ Arg Xaa²¹ Xaa²²⁻³¹ His Asn Xaa³⁴ Term, wherein:

Xaa¹ is Ser or Ala;

Xaa⁷ is Leu or Phe;

Xaa⁸ is Met or Nle;

Xaa¹⁶ is Asn or Ser;

25 Xaa¹⁸ is Leu, Met, or Nle;

Xaa¹⁹ is Glu or Arg;

Xaa²¹ is Val or Arg;

Xaa²²⁻³¹ is selected from (SEQ ID NO:26, 27, 28, 29, and 30);

Xaa³⁴ is Phe or Tyr;

30 Term is OH or NR₂, where each R is H or (C₁-C₄)alkyl; and the pharmaceutically acceptable salts thereof. (Formula II)

Representative polypeptides include, but are not limited to:

35 Ala Val Ser Glu Ile Gln Phe Nle His Asn Leu Gly Lys His Leu
 1 5 10 15
Ser Ser Nle Glu Arg Val Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

40 Leu His Asn Tyr NH₂ (SEQ ID NO:23) and

45 Ala Val Ser Glu Ile Gln Phe Nle His Asn Leu Gly Lys His Leu
 1 5 10 15
Ser Ser Nle Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Asn Tyr NH₂ (SEQ ID NO:24).

50

In still another aspect of this invention, it has surprisingly been found that homologs and analogs of PTH and PTHrP having less than 34 amino acids are also potent bone remodeling agents. These compounds are of

-18-

general formula:

Ala Val Ser Glu Xaa⁵ Gln Leu Leu His Asp Xaa¹¹ Gly Xaa¹³ Ser Ile Gln Asp Leu
Xaa¹⁹ Arg Xaa²¹ Xaa²²⁻³¹ Xaa³² Xaa³³ Xaa³⁴ Term,

5 Representative polypeptides include, but are not limited to:

Compound 41: AVSEHQLLHD KGKSIQDLRR RELLEKLLEK LHP-NH₂ (SEQ ID NO:55)

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile

1 5 10 15
10 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30
Leu His Pro NH₂ (SEQ ID NO:55).

Physical Data

15 m.p. 142.8-166.1°C [α]_D²⁵ -53.80 (c 0.38, H₂O)
FAB (C₁₇₃H₂₈₅N₅₅O₄₉): [M+H]⁺ 3929
AAA: Asx 2.0 (2) Glx 5.7 (6) Ser 1.8 (2)
His 3.0 (3) Gly 1.1 (1) Ala 0.9 (1)
Arg 2.8 (3) Val 1.2 (1) Ile 0.9 (1)
20 Leu 7.4 (8) Lys 4.4 (4) Pro 0.9 (1)

Compound 42: AVSEHQLLHD KGKSIQDLRR RELLEKLLEK LP-NH₂ (SEQ ID NO:56)

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile

1 5 10 15
25 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30
Leu Pro NH₂ (SEQ ID NO:56).

Physical Data

30 m.p. 161.0-177.0°C [α]_D²⁵ -61.97 (c 0.19, H₂O)
FAB (C₁₆₇H₂₈₈N₅₂O₄₈): [M+H]⁺ 3792.0
AAA: Asx 2.2 (2) Glx 5.9 (6) Ser 1.9 (2)
His 2.1 (2) Gly 1.1 (1) Ala 1.0 (1)
Arg 3.0 (3) Val 1.1 (1) Ile 1.0 (1)
35 Leu 7.9 (8) Lys 4.3 (4) Pro 0.9 (1)

The skilled artisan will appreciate that numerous permutations of the polypeptide analogs may be synthesized which will possess the desirable attributes of those described herein provided that an amino acid sequence having a mean hydrophobic moment per residue at 100° ± 20° greater than about 0.20 is inserted at positions (22-31).

Classical Synthesis of the Polypeptides

The polypeptides of the instant invention may be synthesized by methods such as those set forth in

J.M. Stewart and J.D. Young, *Solid Phase Peptide Synthesis*, 2nd ed., Pierce Chemical Co., Rockford, Illinois (1984) and J. Meienhofer, *Hormonal Proteins and Peptides*, Vol. 2, Academic Press, New York, (1973) for solid phase synthesis and E. Schroder and K. Lubke, *The Peptides*, Vol. 1, Academic Press, New York, (1965) for solution synthesis.

In general, these methods involve the sequential addition of protected amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid and any reactive side chain group are protected. This protected amino acid is then either attached to an inert solid support, or utilized in solution, and the next amino acid in the sequence, also suitably protected, is added under conditions amenable to formation of the amide linkage. After all the desired amino acids have been linked in the proper sequence, protecting groups and any solid support are removed to afford the crude polypeptide. The polypeptide is desalted and purified, preferably chromatographically, to yield the final product.

A preferred method of preparing the analogs of the physiologically active truncated polypeptides, having fewer than about forty amino acids, involves solid phase peptide synthesis. In this method the α -amino (N^{α}) functions and any reactive side chains are protected by acid- or base-sensitive groups. The protecting group should be stable to the conditions of peptide linkage formation, while being readily removable without affecting the extant polypeptide chain. Suitable α -amino protecting groups include, but are not limited to *t*-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), *o*-chlorobenzyloxycarbonyl, biphenylisopropylloxycarbonyl, *t*-amyloxycarbonyl (Amoc), isobornyloxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxy-carbonyl, *o*-nitrophenylsulfenyl, 2-cyano-*t*-butoxycarbonyl, 9-fluorenylmethoxycarbonyl (Fmoc) and the like, preferably *t*-butoxycarbonyl (Boc). Suitable side chain protecting groups include, but are not limited to: acetyl, benzyl (Bzl), benzyloxymethyl (Bom), *o*-bromobenzyloxycarbonyl, *t*-butyl, *t*-butyldimethylsilyl, 2-chlorobenzyl (Cl-z), 2,6-dichlorobenzyl, cyclohexyl, cyclopentyl, isopropyl, pivalyl, tetrahydropyran-2-yl, tosyl (Tos), trimethylsilyl, and trityl.

In solid phase synthesis, the C-terminal amino acid is first attached to a suitable resin support. Suitable resin supports are those materials which are inert to the reagents and reaction conditions of the stepwise condensation and deprotection reactions, as well as being insoluble in the media used. Examples of commercially available resins include styrene/divinylbenzene resins modified with a reactive group, e.g., chloromethylated co-poly-(styrene-divinylbenzene), hydroxymethylated co-poly-(styrene-divinylbenzene), and the like. Benzylated, hydroxy-

methyalted phenylacetamidomethyl (PAM) resin is preferred. When the C-terminus of the compound is an amide, a preferred resin is p-methylbenzhydrylamino-co-poly(styrene-divinyl-benzene) resin.

Attachment to the PAM resin may be accomplished by reaction of the N⁺-protected amino acid, preferably the Boc-amino acid, as its ammonium, cesium, triethylammonium, 1,5-diazabicyclo-[5.4.0]undec-5-ene, tetramethylammonium, or similar salt in ethanol, acetonitrile, N,N-dimethylformamide (DMF), and the like, preferably the cesium salt in DMF, with the resin at an elevated temperature, for example between about 40° and 60°C, preferably about 50°C, for from about 12 to 72 hours, preferably about 48 hours.

The N⁺-Boc-amino acid may be attached to the benzhydrylamine resin by means of, for example, an N,N'-diisopropylcarbodiimide (DIC)/1-hydroxybenzotriazole (HOBt) mediated coupling for from about 2 to about 24 hours, preferably about 2 hours at a temperature of between about 10° and 50°C, preferably 25°C in a solvent such as dichloro-methane or dimethylformamide, preferably dichloromethane.

The successive coupling of protected amino acids may be carried out by methods well known in the art, typically in an automated peptide synthesizer. Following neutralization with triethylamine or similar base, each protected amino acid is preferably introduced in approximately 1.5 to 2.5 fold molar excess and the coupling carried out in an inert, nonaqueous, polar solvent such as dichloromethane, DMF, or mixtures thereof, preferably in dichloromethane at ambient temperature. Representative coupling agents are N,N'-dicyclohexylcarbodiimide (DCC), N,N'-diisopropyl-carbodiimide (DIC) or other carbodiimide, either alone or in the presence of 1-hydroxybenzotriazole (HOBt), O-acyl ureas, benzotriazol-1-yl-oxytris(pyrrolidino)phosphonium hexafluorophosphate (PyBop), N-hydroxysuccinimide, other N-hydroxyimides, or oximes. Alternatively, protected amino acid active esters (e.g. p-nitrophenyl, pentafluorophenyl and the like) or symmetrical anhydrides may be used.

At the end of the solid phase synthesis the fully protected peptide is removed from the resin. When the linkage to the resin support is of the benzyl ester type, cleavage may be effected by means of aminolysis with an alkylamine or fluoroalkylamine for peptides with an alkylamide C-terminus, or by aminolysis with, for example, ammonia/methanol or ammonia/ethanol for peptides with an unsubstituted amide C-terminus, at a temperature between about -10° and 50°C, preferably about 25°C, for between about 12 and 24 hours, preferably about 18 hours. Peptides with a hydroxy C-terminus may be cleaved by HF or other strongly acidic deprotection regimen or by saponification. Alternatively, the peptide may be removed from the resin by transesterification, e.g., with methanol, followed by aminolysis or saponification. The protected peptide may be purified by silica gel chromatography.

The side chain protecting groups may be removed from the peptide by

treating the aminolysis product with, for example, anhydrous liquid hydrogen fluoride in the presence of anisole or other carbonium ion scavenger, treatment with hydrogen fluoride/pyridine complex, treatment with tris(trifluoroacetyl)boron and trifluoroacetic acid, by reduction with hydrogen and palladium on carbon or polyvinylpyrrolidone, or by reduction with sodium in liquid ammonia, preferably with liquid hydrogen fluoride and anisole at a temperature between about -10° and +10°C, preferably at about 0°C, for between about 15 minutes and 2 hours, preferably about 1.5 hours.

For peptides on the benzhydrylamine resin, the resin cleavage and deprotection steps may be combined in a single step utilizing liquid hydrogen fluoride and anisole as described above.

The solution may be desalted (e.g. with BioRad AG-3[®] anion exchange resin) and the peptide purified by a sequence of chromatographic steps employing any or all of the following types: ion exchange on a weakly basic resin in the acetate form; hydrophobic adsorption chromatography on underivatized co-poly(styrene-divinylbenzene), e.g. Amberlite[®] XAD; silica gel adsorption chromatography; ion exchange chromatography on carboxymethylcellulose; partition chromatography, e.g. on Sephadex[®] G-25; countercurrent distribution; or high performance liquid chromatography (HPLC), especially reversed-phase HPLC on octyl- or octadecylsilylsilica (ODS) bonded phase column packing.

Thus, another aspect of the present invention relates to processes for preparing polypeptides and pharmaceutically acceptable salts thereof which processes comprise sequentially condensing protected amino acids on a suitable resin support, removing the protecting groups and resin support, and purifying the product, to afford analogs of the physiologically active truncated homologs and analogs of PTH and PTHrp, preferably of PTH(1-34) and PTHrp(1-34), in which the amino acids at positions (22-31) form an amphipathic α -helical peptide sequence, as defined above.

Recombinant Synthesis of the Polypeptides

Alternatively, the polypeptides of this invention may be prepared by cloning and expression of a gene encoding for the desired polypeptide. In this process, a plasmid containing the desired DNA sequence is prepared and inserted into an appropriate host microorganism, typically a bacteria, such as *E. coli*, or a yeast, such as *Saccharomyces cerevisiae*, inducing the host microorganism to produce multiple copies of the plasmid, and so of the cDNA encoding for the polypeptide analogs of this invention.

First, a synthetic gene coding for the selected PTH or PTHrp analog is designed with convenient restriction enzyme cleavage sites to facilitate subsequent alterations. Polymerase chain reaction (PCR), as taught by Mullis in U. S. Patent Nos. 4,683,195 and 4,683,202, may be used to amplify the sequence.

The amplified synthetic gene may be isolated and ligated to a

suitable plasmid, such as a Trp LE plasmid, into which four copies of the gene may be inserted in tandem. Preparation of Trp LE plasmids is described in U.S. Patent No. 4,738,921 and European Patent Publication No. 0212532. Trp LE plasmids generally produce 8-10 times more protein than Trp E plasmids. The multi-copy gene may then be expressed in an appropriate host, such as *E. coli* or *S. cerevisiae*.

The specific expression vector used herein was Trp LE 18 Prot (Ile³, Pro⁵) containing the following elements: a pBR322 fragment (EcoRI-BamHI) containing the ampicillin resistant gene and the plasmid origin of replication; an EcoRI-SacII fragment containing the trp promoter and the trpE gene; an HIV protease (Ile³,Pro⁵) gene fragment (SacII-HindIII); a bGRF gene fragment (HindIII-BamHI); and a transcription terminator from *E. coli* rpoc gene. The HIV protease and bGRF gene fragments are not critical and may be replaced with other coding sequences, if desired.

The expressed multimeric fusion proteins accumulate intracellularly into stable inclusion bodies and may be separated by centrifugation from the rest of the cellular protein. The isolated fusion protein is converted to the monomeric PTH or PTHrp analog and may be purified by cation exchange and/or reverse phase HPLC.

Alternative methods of cloning, amplification, expression, and purification will be apparent to the skilled artisan. Representative methods are disclosed in Maniatis, et al., *Molecular Cloning, a Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory (1989).

Utility and Administration

The polypeptides of this invention are useful for the prevention and treatment of a variety of mammalian conditions manifested by loss of bone mass. In particular, the compounds of this invention are indicated for the prophylaxis and therapeutic treatment of osteoporosis and osteopenia in humans.

In general, the polypeptides of this invention, or salts thereof, are administered in amounts between about 0.002 and 1 $\mu\text{g/kg}$ body weight per day, preferably from about 0.04 to about 0.2 $\mu\text{g/kg}$ body weight per day. For a 50 kg human female subject, the daily dose of active ingredient is from about 0.1 to about 50 μgs , preferably from about 2.0 to about 10 μgs . In other mammals, such as horses, dogs, and cattle, higher doses may be required. This dosage may be delivered in a conventional pharmaceutical composition by a single administration, by multiple applications, or via controlled release, as needed to achieve the most effective results, preferably one or more times daily by injection.

The selection of the exact dose and composition and the most appropriate delivery regimen will be influenced by, *inter alia*, the pharmacological properties of the selected polypeptide, the nature and severity of the condition being treated, and the physical condition and

mental acuity of the recipient.

Representative delivery regimens include oral, parenteral (including subcutaneous, intramuscular and intravenous), rectal, buccal (including sublingual), pulmonary, transdermal, and intranasal.

5 Pharmaceutically acceptable salts retain the desired biological activity of the parent polypeptide without toxic side effects. Examples of such salts are (a) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; and salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, 10 maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, pamoic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acids, naphthalene disulfonic acids, polygalacturonic acid and the like; (b) base addition salts formed with 15 polyvalent metal cations such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium, and the like; or with an organic cation formed from N,N'-dibenzylethylenediamine or ethylenediamine; or (c) combinations of (a) and (b), e.g., a zinc tannate salt and the like.

20 A further aspect of the present invention relates to pharmaceutical compositions comprising as an active ingredient a polypeptide of the present invention, or pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable, non-toxic carrier. As mentioned above, such compositions may be prepared for parenteral (subcutaneous, intramuscular or intravenous) administration, particularly 25 in the form of liquid solutions or suspensions; for oral or buccal administration, particularly in the form of tablets or capsules; for pulmonary or intranasal administration, particularly in the form of powders, nasal drops or aerosols; and for rectal or transdermal administration.

30 The compositions may conveniently be administered in unit dosage form and may be prepared by any of the methods well-known in the pharmaceutical art, for example as described in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA., (1985). Formulations for parenteral administration may contain as excipients sterile water or 35 saline, alkylene glycols such as propylene glycol, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes and the like. For oral administration, the formulation can be enhanced by the addition of bile salts or acylcarnitines. Formulations for nasal administration may be solid and may contain excipients, for example, 40 lactose or dextran, or may be aqueous or oily solutions for use in the form of nasal drops or metered spray. For buccal administration typical excipients include sugars, calcium stearate, magnesium stearate, pregelatinated starch, and the like.

45 When formulated for nasal administration, the absorption across the nasal mucous membrane may be enhanced by surfactant acids, such as for

example, glycocholic acid, cholic acid, taurocholic acid, ethocholic acid, deoxycholic acid, chenodeoxycholic acid, dehydrocholic acid, glycodeoxycholic acid, cyclodextrins and the like in an amount in the range between about 0.2 and 15 weight percent, preferably between about 0.5 and 4 weight percent, most preferably about 2 weight percent.

Delivery of the compounds of the present invention to the subject over prolonged periods of time, for example, for periods of one week to one year, may be accomplished by a single administration of a controlled release system containing sufficient active ingredient for the desired release period. Various controlled release systems, such as monolithic or reservoir-type microcapsules, depot implants, osmotic pumps, vesicles, micelles, liposomes, transdermal patches, iontophoretic devices and alternative injectable dosage forms may be utilized for this purpose. Localization at the site to which delivery of the active ingredient is desired is an additional feature of some controlled release devices, which may prove beneficial in the treatment of certain disorders.

One form of controlled release formulation contains the polypeptide or its salt dispersed or encapsulated in a slowly degrading, non-toxic, non-antigenic polymer such as copoly(lactic/glycolic) acid, as described in the pioneering work of Kent, Lewis, Sanders, and Tice, U.S. 4,675,189. The compounds or, preferably, their relatively insoluble salts, may also be formulated in cholesterol or other lipid matrix pellets, or silastomer matrix implants. Additional slow release, depot implant or injectable formulations will be apparent to the skilled artisan. See, for example, *Sustained and Controlled Release Drug Delivery Systems*, J. R. Robinson ed., Marcel Dekker, Inc., New York, 1978, and R. W. Baker, *Controlled Release of Biologically Active Agents*, John Wiley & Sons, New York, 1987.

The following specific Examples are intended to illustrate the synthesis and testing of representative compounds of the invention and should not be construed as limiting the scope of the claims. In the Examples, "m. p." is melting point, " $[\alpha]_D^{25}$ " is the optical activity at 25°C at the given concentration in the indicated solvent, "FAB" is fast atom bombardment mass spectrometry, and "AAA" is amino acid analysis, with expected values in parentheses following the observed values. The amino acid analyses were conducted on a Hewlett-Packard AminoQuant Analyzer following the manufacturer's recommended protocols. Primary amino acids were derivatized with o-phthalaldehyde; secondary amino acids with Fmoc. Fluorescent detection of the derivatized amino acids was used for quantification. The protected amino acids were obtained from Applied Biosystems Inc. (Foster City, CA).

EXAMPLE I

Compound 1 (SEQ ID NO:7) was prepared on 0.5 mmol scale by the solid phase method on 4-methylbenzhydrylamine resin, using an automated Applied Biosystems Model 430A peptide synthesizer. The α -amino groups were protected with t-butoxycarbonyl (Boc). The side chain protecting groups were: benzyl (Bzl) for Asp, Glu, and Ser; tosyl (Tos) for Arg; benzyloxymethyl (Bom) for His; and 2-chlorobenzyl (Cl-2) for Lys. The amino acids were coupled sequentially using N,N-dicyclohexylcarbodiimide/1-hydroxybenzotriazole (DCC/HOBt) following Stewart and Young (*supra*). After each amino acid coupling, the peptide was acetylated using a mixture of acetic anhydride and diisopropylethylamine in N-methylpyrrolidone. The completed peptide was cleaved from the resin with simultaneous deprotection of the side chain protecting groups using anhydrous HF (25 mL) in the presence of anisole (2.5 mL) at -10°C for 30 minutes and at 0°C for 60 minutes. After evaporation of the HF in vacuo, the residue was washed with anhydrous ether, and the crude peptide extracted with 10% acetic acid. Lyophilization of the 10% acetic acid extract gave 900 mgs of crude product. The peptide was purified by medium pressure ODS reversed phase column chromatography using a gradient of 22-45% CH₃CN in 0.1% TFA. The product eluted in three fractions, which were concentrated and lyophilized to give 130 mgs of white solid of >98% purity.

Compound 1:

AVSEHOLLHDKGKSIODLRRLLEKLLEKLHTA-NH₂ (SEQ ID NO:7)

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Thr Ala NH₂ (SEQ ID NO:7)

Physical Data: m.p. 150-159°C [α]_D²⁵ -34.88° (c 0.16, H₂O)

FAB (C₁₇₃H₃₀₀N₅₆O₅₁): [M+H]⁺ 4005.5

AAA: Asp, 1.9(2); Glu, 5.6(6); Ser, 1.6(2); His, 2.7(3); Gly, 1.0(1); Thr, 0.9(1); Ala, 1.9(2); Arg, 2.8(3); Val, 1.0(1); Ile, 0.9(1); Leu, 7.3(8); Lys, 4.0(4).

Similarly, Compounds 2, 5-18, 21-27, 29-36, 38-48, 50-54, 58-64 and 66-70 were prepared and characterized, substituting PAM resin as needed for the synthesis of hydroxy-terminal polypeptides.

AVSEHOLLHDKGKSIQDLRRRELLEKLLEKLHTA-OH (SEQ ID NO:6)

Physical Data: m.p. 154-170°C $[\alpha]_D^{25}$ -49.35° (c 0.46, H₂O)
 FAB (C₁₇₅H₃₀₁N₅₇O₅₀): [M+H]⁺ 4005.0
 AAA: Asp, 2.1(2); Glu, 5.9(6); Ser, 1.7(2); His, 2.9(3); Gly 1.1(1); Thr,
 1.0(1); Ala, 1.9(2); Arg, 3.0(3); Val, 1.2(1); Ile, 1.0(1); Leu,
 7.8(8); Lys, 4.2(4).

AVSEHOLLHDKGKSIODLRRRELLERLLERLHTA-OH (SEQ ID NO:15)

Physical Data: m.p. 147-165°C $[\alpha]_D^{25}$ -49.17° (c 0.66, H₂O)
FAB (C₁₇₅H₂₉₉N₅₃O₅₂): [M+H]⁺ 4061
AAA: Asp, 2.1(2); Gly, 6.1(6); Ser, 1.8(2); His, 3.1(3); Gly, 1.1(1); Thr,
1.0(1); Ala, 2.0(2); Arg, 5.0(5); Val, 1.0(1); Ile, 0.9(1); Leu,
7.7(8); Lys, 1.9(2).

AVSEHOLLHDRGRSIODLRRRELLERLLERLHTA-OH (SEQ ID NO:16)

Physical Data: m.p. 150-170°C $[\alpha]_D^{25}$ -48.65° (c 0.54, H₂O)
FAB (C₁₇₅H₂₉₉N₆₃S₅₂): [M+H]⁺ 4118.0
AAA: Asp, 2.1(2); Glu, 6.1(6); Ser, 1.8(2); His, 3.2(3); Gly, 1.2(1); Thr,
1.0(1); Ala, 2.0(2); Arg, 6.9(7); Val, 1.0(1); Ile, 1.0(1); Leu,
7.8(8).

-27-

Compound 7:AVSEHOLLHDRGRSIODLRRRELLERLLKRLHTA-OH (SEQ ID NO:17)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Lys Arg
 20 25 30
 10 Leu His Thr Ala OH (SEQ ID NO:17)

Physical Data: m.p. 177-182°C $[\alpha]_D^{25}$ -46.17° (c 0.14, H₂O)
 FAB (C₁₇₆H₃₀₄N₆₄O₅₀): [M+H]⁺ 4117
 15 AAA: Asp, 2.0(2); Glu, 4.8(5); Ser, 1.8(2); His, 3.2(3); Gly, 1.1(1); Thr,
 0.9(1); Ala, 1.9(2); Arg, 6.7(7); Val, 1.0(1); Ile, 1.0(1); Lys,
 7.7(8); Lys, 0.9(1).

20

Compound 8:AVSEHOLLHDKGKSIODLRRRELLEKLLRKLHTA-OH (SEQ ID NO:5)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Arg Lys
 20 25 30
 30 Leu His Thr Ala OH (SEQ ID NO:5)

Physical Data: m.p. 147-165°C $[\alpha]_D^{25}$ -49.17° (c 0.66, H₂O)
 FAB (C₁₇₆H₃₀₄N₅₉O₄₉): [M+H]⁺ 4033.0
 35 AAA: Asp, 2.0(2); Glu, 4.8(5); Ser, 1.8(2); His, 2.7(3); Gly, 1.1(1); Thr,
 0.9(1); Ala, 2.0(2); Arg, 3.9(4); Val, 1.0(1); Ile, 1.0(1); Leu,
 7.9(8); Lys, 4.0(4).

40

Compound 9:AVSEHOLLHDKGKSIODLRRRELLEKLLKLTAGRR-OH (SEQ ID NO:10)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 50 Leu His Thr Ala Gly Arg Arg OH (SEQ ID NO:10)
 35

Physical Data: m.p. 158-160°C $[\alpha]_D^{25}$ -44.76° (c 0.1, H₂O)
 FAB (C₁₈₉H₃₂₆N₆₄O₅₅): [M]⁺ 4375.0
 55 AAA: Asp, 2.0(2); Glu, 5.9(6); Ser, 1.7(2); His, 2.9(3); Gly, 2.3(2); Thr,
 1.0(1); Ala, 1.9(2); Arg, 5.0(5); Val, 1.2(1); Ile, 1.0(1); Leu,
 7.8(8); Lys, 4.3(4).

60

AVSEAOLLHDLGKSIODLRRRELLEKLEKLHAL-OH (SEQ ID NO:14)

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Ala Leu OH (SEQ ID NO:14)

15 Physical Data: m.p. 170-175°C $[\alpha]_D^{25}$ -31.59° (c 0.54, H₂O)
FAB (C₁₇₄H₃₀₀N₅₂O₅₁): [M+H]⁺ 3936.0
AAA: Asp, 2.0(2); Glu, 6.0(6); Ser, 1.8(2); His, 2.0(2); Gly, 1.2(1); Ala,
3.0(3); Arg, 2.8(3); Val, 1.1(1); Ile, 1.0(1); Leu, 9.9(10); Lys,
3.0(3).

AVSEHOLLHDKGKSIQDLRRELLEKLLLELLKEL-NH, (SEQ ID NO:11)

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu Lys Glu Leu NH₂ (SEQ ID NO:11)

35 Physical Data: m.p. 172-174°C $[\alpha]_D^{25}$ -43.29° (c 0.2, H₂O)
FAB (C₁₇₉H₃₁₁N₅₅O₅₂): [M+H]⁺ 4065.8
AAA: Asp, 2.2(2); Glu, 7.7(7); Ser, 1.7(2); His, 2.0(2); Gly, 1.0(1); Ala,
1.0(1); Arg, 3.0(3); Val, 1.1(1); Ile, 1.0(1); Leu, 9.3(9); Lys,
5.1(5).

AVSEIQFXHNLGKHLSSXERVELLEKLLEKLHNY-NH. (X=N1e, SEQ ID NO:23)

Ser Ser Nle Glu Arg Val Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

50 Leu His Asn Tyr NH₂ (SEQ ID NO:23)

55 Physical Data: m.p. 178°C [α]_D²⁵ -36.88° (c 0.4, H₂O)
FAB (C₁₈₂H₂₉₉N₅₁O₅₁): [M+H]⁺ 4001.6
AAA: Asp, 2.1(2); Glu, 6.5(6); Ser, 2.7(3); His, 3.1(3); Gly, 1.1(1); Ala,
1.0(1); Arg, 1.0(1); Tyr, 0.8(1); Val, 2.0(2); Phe, 1.0(1); Ile,
0.9(1); Leu+Nle, 8.5(7+2); Lys, 3.1(3).

```

5      Ala Val Ser Glu Ile Gln Phe Nle His Asn Leu Gly Lys His Leu
      1          5          10          15

      Ser Ser Nle Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
      20          25          30

10     Leu His Asn Tyr NH2 (SEQ ID NO:24)

```

FAB (C₁₈₄H₃₀₄N₅₆O₄₉) : [M+H]⁺ 4084

20 Compound 14:

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15
Gln Asp Leu Arg Arg Arg Ala Leu Ala Glu Ala Leu Ala Glu Ala
20 25 30

FAB (C₁₆H₂₇N₃O₄) : [M+H]⁺ 3747.0

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
1 5 10 15
Gln Asp Leu Ala Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

FAB ($C_{177}H_{293}N_{53}O_{51}$): $[M+H]^+$ 3919.0

BNSDOCID: <WO__9401460A1_I_>

Compound 16:

60

-31-

Compound 21:AVSEIOFLHN LGKHLSSLRR RELLEKLLEK LHNY-NH₂ (SEQ ID NO:35)

5 Ala Val Ser Glu Ile Gln Phe Leu His Asn Leu Gly Lys His Leu
 1 5 10 15
 Ser Ser Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 10 Leu His Asn Tyr NH₂ (SEQ ID NO:35)

Physical Data: m.p. 148-155°C $[\alpha]_D^{25}$ -45.97 (c 0.26, H₂O)

FAB (C₁₈₄H₃₀₄N₅₆O₄₉): [M+H]⁺ 4084

15 AAA: Asx, 2.1(2); Glx, 5.0(5); Ser, 2.7(3); His, 3.0(3); Gly, 1.0(1); Ala,
 0.9(1); Arg, 3.1(3); Tyr, 0.9(1); Val, 1.0(1); Phe, 0.9(1); Ile,
 0.9(1); Leu 9.3(9); Lys, 3.2(3).

20

Compound 24:AVSEHOLLHD KGKSIODLKL KELLEKLLEK LHNTA-NH₂ (SEQ ID NO:38)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Lys Leu Lys Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 30 Leu His Thr Ala NH₂ (SEQ ID NO:38)

Physical Data: m.p. 175-182°C $[\alpha]_D^{25}$ -49.99 (c 0.47, H₂O)

FAB (C₁₇₅H₂₉₉N₄₉O₅₁): [M+H]⁺ 3906.5

35 AAA: Asx, 2.1(2); Glx, 6.5(6); Ser, 1.8(2); His, 3.1(3); Gly, 1.1(1); Thr,
 1.0(1); Ala, 2.1(2); Val, 1.1(1); Ile, 1.0(1); Leu, 9.1(9); Lys,
 6.5(6).

40

Compound 25:AVSEHOLLHD KGKSIODLRR RELLERLLER LHNTA-NH₂ (SEQ ID NO:39)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30

50

Leu His Thr Ala-NH₂ (SEQ ID NO:39)

Physical Data: m.p. 136.5-153.5°C $[\alpha]_D^{25}$ -32.57 (c 0.13, H₂O)

FAB (C₁₇₅H₃₀₀N₅₀O₅₁): [M+H]⁺ 4060.8

55 AAA: Asx, 2.2(2); Glx, 6.2(6); Ser, 1.8(2); His, 3.2(3); Gly, 1.1(1); Thr,
 1.0(1); Ala, 2.1(2); Arg, 5.2(5); Val, 1.1(1); Ile, 1.1(1); Leu,
 8.4(8); Lys, 2.2(2).

60

-32-

Compound 26:AVSEHOLLHD KGKSIODLRR RELLERLLER LHTAP-OH (SEQ ID NO:40)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30

10 Leu His Thr Ala Pro OH (SEQ ID NO:40)

Physical Data: m.p. 125.8-127.2°C $[\alpha]_D^{25}$ -54.62(c 0.23, H₂O)
 FAB(C₁₈₀H₃₀₆N₆₀O₅₃): [M+H]⁺ 4158.0
 15 AAA: Asx, 2.1(2); Glx, 6.2(6); Ser, 1.8(2); His, 2.9(3); Gly, 1.1(1); Thr,
 1.0(1); Ala, 2.0(2); Arg, 5.1(5); Val, 1.0(1); Ile, 1.0(1); Leu,
 8.0(8); Lys, 2.1(2); Pro, 1.1(1).

20

Compound 27:AVSEHOLLHD KGKSIODLRR RELLERLLER LHTAGRR-OH (SEQ ID NO:41)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30

30 Leu His Thr Ala Gly Arg Arg OH (SEQ ID NO:41)
 35

Physical Data: m.p. 106-137.3°C $[\alpha]_D^{25}$ -39.55(c 0.67, H₂O)
 35 FAB(C₁₈₉H₃₂₆N₆₈O₅₃): [M+H]⁺ 4430.5
 AAA: Asx, 2.1(2); Glx, 5.9(6); Ser, 1.6(2); His, 2.7(3); Gly, 2.2(2); Thr,
 1.0(1); Ala, 1.8(2); Arg, 7.3(7); Val, 0.8(1); Ile, 1.0(1); Leu,
 8.1(8); Lys, 2.1(2).

40

Compound 29:AVSEHOLLHD KGKSIODLRR RELLEKLEK LHTY-NH₂ (SEQ ID NO:43)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15

50 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Thr Tyr NH₂ (SEQ ID NO:43)

Physical Data: m.p. 160-172°C $[\alpha]_D^{25}$ -49.85(c 0.34, H₂O)
 55 FAB(C₁₈₁H₃₀₄N₅₆O₅₂): [M+H]⁺ 4096.9
 AAA: Asx, 2.0(2); Glx, 5.6(6); Ser, 1.7(2); His, 3.1(3); Gly, 1.1(1); Thr,
 0.9(1); Ala, 0.9(1); Arg, 3.0(3); Tyr, 0.9(1); Val, 1.0(1); Ile,
 1.0(1); Leu, 7.7(8); Lys, 4.4(4).

60

Compound 30:

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Tyr Ser Ile
1 5 10 15
Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30
10 Leu His Thr Ala NH. (SEQ ID NO:44)

15 AAA: Asx, 2.0(2); Glx, 5.5(6); Ser, 1.8(2); His, 3.4(3); Gly, 1.1(1); Thr, 1.0(1); Ala, 2.0(2); Arg, 2.9(3); Tyr, 0.8(1); Val, 1.0(1); Ile, 0.9(1); Leu, 7.9(8); Lys, 3.4(3).

Compound 31:

```

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Cys Ser Ile
   1      5      10
   Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
   20      25      30
30 Leu His Thr Ala NH, (SEQ ID NO:45)

```

35 AAA: Asx+Cys, 3.0(2+1); Glx, 5.6(6); Ser, 1.7(2); His, 3.0(3); Gly,
1.0(1); Thr, 0.9(1); Ala, 1.9(2); Arg, 2.5(3); Val, 1.0(1); Ile,
0.9(1); Leu, 7.5(8); Lys, 3.3(3).

Compound 32:

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Xaa Ser Ile
1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
20 25 30

Leu His Thr Ala NH₂, (Xaa = Cys(CH₂CONH(CH₂)₃NH(biotinyl))), (SEQ ID NO:46)

AAA: Asx, 2.2(2); Glx, 6.1(6); Ser, 1.8(2); His, 3.8(3); Gly, 1.0(1); Thr, 1.0(1); Ala, 2.0(2); Arg, 3.1(3); Val, 1.1(1); Ile, 0.9(1); Leu, 8.3(3); Lys, 3.3(3).

-34-

Compound 33:

AVSEHOLLHD KGXSIQDLRR RELLEKI LEK LH7A-NH₂ (SEQ ID NO:47)
(X = Lys(7-dimethylamino-2-oxo-2H-1-benzopyran-4-acetyl))

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Xaa Ser Ile
 1 5 10 15
 10 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Thr Ala NH₂, (Xaa = Lys(7-dimethylamino-2-oxo-2H-1-benzopyran-4-acetyl), (SEQ ID NO:47)

15 Physical Data: m.p. 135-205°C $[\alpha]_D^{25}$ -26.92 (c 0.104, 50% aq. HOAc)
 FAB(C₁₈₈H₃₁₁N₅₇O₅₄): [M+H]⁺ 4233
 20 AAA: Asx, 1.9(2); Glx, 6.3(6); Ser, 1.7(2); His, 3.2(3); Gly, 1.0(1); Thr, 1.1(1); Ala, 2.0(2); Arg, 3.2(3); Val, 1.1(1); Ile, 0.9(1); Leu, 8.2(8); Lys, 4.5(4).

Compound 34:

AVSEHOLLHD KGKSIQDLRR RELLEKLLEK LH7AG-OH (SEQ ID NO:48)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 30 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

35 Leu His Thr Ala Gly OH (SEQ ID NO:48)
 35

Physical Data: m.p. 92.1-146.6°C $[\alpha]_D^{25}$ -40.76 (c 0.34, H₂O)
 FAB(C₁₇₇H₃₀₂N₅₆O₅₃): [M+H]⁺ 4062.0
 40 AAA: Asx, 2.0(2); Glx, 5.7(6); Ser, 1.8(2); His, 3.0(3); Gly, 2.2(2); Thr, 0.9(1); Ala, 1.9(2); Arg, 2.8(3); Val, 1.2(1); Ile, 0.9(1); Leu, 7.5(8); Lys, 4.2(4).

Compound 35:

AVSX₁HOLLHX₂ KGKSIOX₁LRR RX₁LLX₂KLLX₁K LHA-OH (SEQ ID NO:49)
(X₁ = Glu(OCH₃); X₂ = Asp(OCH₃))

50 Ala Val Ser Xaa₁ His Gln Leu Leu His Xaa₂ Lys Gly Lys Ser Ile
 1 5 10 15
 55 Gln Xaa₁ Leu Arg Arg Arg Xaa₁ Leu Leu Xaa₁ Lys Leu Leu Xaa₁ Lys
 20 25 30

Leu His Ala OH (Xaa₁ = Glu(OCH₃); Xaa₂ = Asp(OCH₃)) (SEQ ID NO:49)

Physical Data: m.p. (not determined) $[\alpha]_D^{25}$ -21.96 (c 0.132, H₂O)
 FAB(C₁₈₁H₃₁₁N₅₅O₅₂): [M+H]⁺ 4089.0
 60 AAA: Asx, 2.1(2); Glx, 6.3(6); Ser, 1.8(2); His, 3.3(3); Gly, 1.1(1); Thr, 1.0(1); Ala, 2.0(2); Arg, 3.1(3); Val, 1.1(1); Ile, 0.9(1); Leu, 8.0(8); Lys, 4.2(4).

65

-35-

Compound 36:

AVSX,HOLLHX, KGKSIOX,LRR RX,LLX,KLLX,K LHA-OCH₃ (SEQ ID NO:50)
(X₁ = Glu(OCH₃); X₂ = Asp(OCH₃))

5 Ala Val Ser Xaa₁ His Gln Leu Leu His Xaa₂ Lys Gly Lys Ser Ile
 1 5 10 15
 10 Gln Xaa₂ Leu Arg Arg Arg Xaa₁ Leu Leu Xaa₁ Lys Leu Leu Xaa₁ Lys
 20 25 30

Leu His Ala OCH₃ (Xaa₁ = Glu(OCH₃); Xaa₂ = Asp(OCH₃)) (SEQ ID NO:50)

15 Physical Data: m.p. (not determined) $[\alpha]_D^{25}$ -46.80 (c 0.07, H₂O)
 FAB(C₁₈₂H₃₁₃N₅₅O₅₂): [M+H]⁺ 4103
 AAA: Asx, 2.1(2); Glx, 6.2(6); Ser, 1.4(2); His, 3.0(3); Gly, 1.1(1); Thr,
 1.1(1); Ala, 1.7(2); Arg, 3.2(3); Val, 0.6(1); Ile, 0.9(1); Leu,
 8.0(8); Lys, 4.1(4).

Compound 38:

AVSEHOLLHD KGKSIODLRR RELLEKLEK LHTAP-OH (SEQ ID NO:52)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 30 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Thr Ala Pro OH (SEQ ID NO:52)
 35

35 Physical Data: m.p. 152.1-186.5°C $[\alpha]_D^{25}$ -55.91 (c 0.33, H₂O)
 FAB(C₁₈₀H₃₀₆N₅₆O₅₃): [M+H]⁺ 4102.6
 AAA: Asx, 2.0(2); Glx, 5.6(6); Ser, 1.7(2); His, 2.9(3); Gly, 1.1(1); Thr,
 0.9(1); Ala, 1.9(2); Arg, 2.9(3); Val, 1.2(1); Ile, 1.0(1); Leu,
 7.7(8); Lys, 4.3(4).

Compound 39:

AVSEHOLLHD KGKSIODLRR RELLEKLEK LHTP-OH (SEQ ID NO:53)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 50 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Thr Pro OH (SEQ ID NO:53)

55 Physical Data: m.p. 120-148.2°C $[\alpha]_D^{25}$ -52.78 (c 0.47, H₂O)
 FAB(C₁₇₇H₃₀₁N₅₅O₅₂): [M+H]⁺ 4031.0
 AAA: Asx, 2.0(2); Glx, 5.5(6); Ser, 1.8(2); His, 2.9(3); Gly, 1.0(1); Thr,
 1.0(1); Ala, 0.9(1); Arg, 2.9(3); Val, 1.2(1); Ile, 0.9(1); Leu,
 7.5(8); Lys, 3.6(3); Pro, 0.9(1).

60

- 36 -

AVSEHOLLHD KGKSIODLRR RELLEKLEK LHTP-NH. (SEQ ID NO:54)

Physical Data: m.p. 133.9-155.1°C $[\alpha]_D^{25}$ -54.22 (c 0.37, H₂O)
FAB (C₁₇₇H₃₀₂N₅₆O₅₁): [M+H]⁺ 4030.7
AAA: Asx, 2.0(2); Glx, 5.6(6); Ser, 1.9(2); His, 2.9(3); Gly, 1.1(1); Thr,
0.9(1); Ala, 0.9(1); Arg, 2.8(3); Val, 1.2(1); Ile, 1.1(1); Leu,
7.8(8); Lys, 4.2(4); Pro, 0.9(1).

AVSEHOLLHD KGKSIQDLRR RELLEKLEK LHP-NH, (SEQ ID NO:55)

Physical Data: m.p. 142.8-166.1°C $[\alpha]_D^{25}$ -53.80 (c 0.38, H₂O)
 FAB(C₁₇₃H₂₉₅N₅₅O₄₉): [M+H]⁺ 3929
 AAA: Asx, 2.0(2); Glx, 5.7(6); Ser, 1.8(2); His, 3.0(3); Gly, 1.1(1); Ala,
 0.9(1); Arg, 2.8(3); Val, 1.2(1); Ile, 0.9(1); Leu, 7.4(8); Lys,
 4.4(4); Pro, 0.9(1).

AVSEHOLLHD KGKSIQDLRR RELLEKLEK LP-NH, (SEQ ID NO:56)

Physical Data: m.p. 161.0-177.0°C $[\alpha]_D^{25}$ -61.97 (c 0.19, H₂O)
FAB (C₁₆₇H₂₈₈N₅₂O₄₈): [M+H]⁺ 3792.0
AAA: Asx, 2.2(2); Glx, 5.9(6); Ser, 1.9(2); His, 2.1(2); Gly, 1.1(1); Ala, 1.0(1); Arg, 3.0(3); Val, 1.1(1); Ile, 1.0(1); Leu, 7.9(8); Lys, 4.3(4); Pro, 0.9(1).

-37-

Compound 43:AVSEHOLLHD KGKSIODLRR RELLEKLEK LHTRSAW-OH (SEQ ID NO:57)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 10 Leu His Thr Arg Ser Ala Trp OH (SEQ ID NO:57)
 35

Physical Data: m.p. 181-202°C $[\alpha]_D^{25}$ -45.14 (c 0.19, H₂O)
 15 FAB(C₁₉₃H₃₂₆N₆₂O₅₆): [M+H]⁺ 4435.2
 AAA: Asx, 2.0(2); Glx, 5.8(6); Ser, 2.8(3); His, 2.8(3); Gly, 1.1(1); Thr,
 0.9(1); Ala, 1.9(2); Arg, 3.7(4); Ile, 0.9(1); Leu, 7.5(8); Lys,
 4.3(4); Trp, 0.9(1).

20

Compound 44:AVSEHOLLHD RGRSIODLRR RELLERLLER LHTRGRTRSAW-OH (SEQ ID NO:58)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 30 20 25 30
 Leu His Thr Arg Gly Arg Arg Thr Arg Ser Ala Trp OH (SEQ ID NO:58)
 35 40

35 Physical Data: m.p. 130-132.2°C $[\alpha]_D^{25}$ -46.66 (c 0.195, H₂O)
 FAB(C₂₁₆H₃₆₅N₈₁O₆₂): [M+H]⁺ 5088.8
 AAA: Asx, 2.2(2); Glx, 6.0(6); Ser, 2.7(3); His, 3.0(3); Gly, 2.2(2); Thr,
 2.1(2); Ala, 3.0(3); Arg, 10.5(10); Val, 0.9(1); Ile, 1.0(1); Leu,
 8.2(8); Trp, 1.0(1).

40

Compound 45:AVSEHOLLHD RGRSIODLRR RELLERLLER LHTRGRTRSAW-NH₂ (SEQ ID NO:59)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 50 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30
 Leu His Thr Arg Gly Arg Arg Thr Arg Ser Ala Trp NH₂ (SEQ ID NO:59)
 35 40

55 Physical Data: m.p. 158-174°C $[\alpha]_D^{25}$ -43.57 (c 0.53, H₂O)
 FAB(C₂₁₆H₃₆₆N₈₂O₆₁): [M+H]⁺ 5087.4
 AAA: Asx, 1.9(2); Glx, 5.6(6); Ser, 2.6(3); His, 3.3(3); Gly, 2.1(2); Thr,
 2.0(2); Ala, 2.9(3); Arg, 10.1(10); Val, 0.9(1); Ile, 1.0(1); Leu,
 8.3(8); Trp 1.1(1).

60

-38-

Compound 46:

AVSEHQLLHD RGXSIODLRR RELLERLER LHTAGRRTRSAW-OH (SEQ ID NO:60)
 (X = Lys(dihydrocinnamoyl))

5 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Xaa Ser Ile
 1 5 10 15
 10 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30
 Leu His Thr Arg Gly Arg Arg Thr Arg Ser Ala Trp OH (Xaa =
 35 40
 15 Lys(dihydrocinnamoyl), (SEQ ID NO:60)

Physical Data: m.p. 165.4-175.2°C $[\alpha]_D^{25}$ -40.43 (c 0.20, H₂O)
 FAB(C₂₂H₃₇N₈O₆): [M+H]⁺ 5191

20 AAA: Asx, 2.1(2); Glx, 6.3(6); Ser, 2.8(3); His, 3.2(3); Gly, 2.1(2); Thr,
 2.0(2); Ala, 3.2(3); Arg, 9.9(9); Val, 1.0(1); Ile, 0.9(1); Leu,
 8.6(8); Lys, 1.1(1); Trp, 1.1(1).

Compound 47:

AVSEIOFXHN LGKHLSSXTR SAWLRKKLOD VHNY-NH. (SEQ ID NO:61)
 (X = norleucine)

30 Ala Val Ser Glu Ile Gln Phe Nle His Asn Leu Gly Lys His Leu
 1 5 10 15
 Ser Ser Nle Thr Arg Ser Ala Trp Leu Arg Lys Lys Leu Gln Asp
 20 25 30
 35 Val His Asn Tyr NH₂ (SEQ ID NO:61)

Physical Data: m.p. 140-160°C $[\alpha]_D^{25}$ -56.88 (c 0.16, H₂O)
 FAB(C₁₈₀H₂₈₇N₅₅O₅₈): [M+H]⁺ 3989.8

40 AAA: Asx, 3.0(3); Glx, 2.9(3); Ser, 3.7(4); His, 2.8(3); Gly, 1.1(1); Thr,
 0.9(1); Ala, 1.9(2); Arg, 2.0(2); Tyr, 1.0(1); Val, 1.7(2); Phe,
 0.9(1); Ile, 0.9(1); Leu+Nle 5.8(2+4); Lys, 3.4(3); Trp, 1.1(1).

Compound 48:

AVSEHQLLHD KGKSIODLRR RELLEKLLEK LHTMA-NH. (SEQ ID NO:62)

50 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 55 Leu His Thr Met Ala NH₂ (SEQ ID NO:62)
 35

Physical Data: m.p. 140-210°C $[\alpha]_D^{25}$ -47.75 (c 0.178, H₂O)
 FAB(C₁₈₀H₃₀₉N₅₇O₅₂S₁): [M+H]⁺ 4135.0

60 AAA: Asx, 2.3(2); Glx, 6.6(6); Ser, 1.4(2); His, 3.2(3); Gly, 1.1(1); Thr,
 1.0(1); Ala, 2.0(2); Arg, 3.1(3); Val, 0.9(1); Met, 1.1(1); Ile,
 1.0(1); Leu, 8.8(8); Lys, 4.4(4).

-39-

Compound 50:AVSEHOLLHD KGKSIODLRR RFFLEKLLEK LHTA-NH₂ (SEQ ID NO:64)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Phe Phe Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 10 Leu His Thr Ala NH₂ (SEQ ID NO:64)
 35

15 Physical Data: m.p. 136.5-156.8°C [α]_D²⁵ -49.89 (c 0.24, H₂O)
 FAB(C₁₈₂H₃₀₀N₅₆O₄₉): [M+H]⁺ 4056.0
 AAA: Asx, 2.2(2); Glx, 5.0(5); Ser, 1.9(2); His, 3.3(3); Gly, 1.0(1); Thr,
 1.0(1); Ala, 2.1(2); Arg, 3.1(3); Val, 1.0(1); Phe, 2.0(2); Ile,
 0.9(1); Leu, 7.2(7); Lys, 3.5(4).

20

Compound 51:AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTA-NH₂ (SEQ ID NO:65)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu His Lys Leu Leu Glu Lys
 20 25 30
 30 Leu His Thr Ala NH₂ (SEQ ID NO:65)

35 Physical Data: m.p. 80.7-141.0°C [α]_D²⁵ -55.38 (c 0.23, H₂O)
 FAB(C₁₇₆H₃₀₀N₅₈O₄₉): [M+H]⁺ 4012.8
 AAA: Asx, 2.2(2); Glx, 4.9(5); Ser, 1.8(2); His, 4.3(4); Gly, 1.1(1); Thr,
 1.0(1); Ala, 2.0(2); Arg, 3.1(3); Val, 1.1(1); Ile, 1.0(1); Leu,
 8.1(8); Lys, 3.9(4).

40

Compound 52:AVSEHOLLHD KGKSIODLRR RELLEHLLEK LHTA-NH₂ (SEQ ID NO:66)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu His Leu Leu Glu Lys
 20 25 30
 50 Leu His Thr Ala NH₂ (SEQ ID NO:66)

55 Physical Data: m.p. 134.3-157.9°C [α]_D²⁵ -50.72 (c 0.45, H₂O)
 FAB(C₁₇₅H₂₉₉N₅₇O₅₁): [M+H]⁺ 4012.8
 AAA: Asx, 2.1(2); Glx, 5.9(6); Ser, 1.8(2); His, 4.2(4); Gly, 1.1(1); Thr,
 1.0(1); Ala, 2.0(2); Arg, 3.0(3); Val, 1.1(1); Ile, 0.9(1); Leu,
 8.1(8); Lys, 3.1(3).

60

AVSEHOLLHD KGKSIODLRR RELLEKLIAK LH7A-NH, (SEQ ID NO:67)

Physical Data: m.p. 142.7-159.8°C $[\alpha]_D^{25}$ -54.01 (c 0.21, H₂O)
FAB (C₁₇₃H₂₉₈N₅₆O₄₉): [M+H]⁺ 3946.0
AAA: Asx, 2.2(2); Glx, 4.9(5); Ser, 1.8(2); His, 3.1(3); Gly, 1.1(1); Thr,
 1.0(1); Ala, 3.1(3); Arg, 3.1(3); Val, 1.0(1); Ile, 1.9(2); Leu,
 7.0(7); Lys, 4.3(4).

AVSEHOLLHD KGKSIODLRR RELLEKLEE IHTA-NH, (SEQ ID NO:68)

Physical Data: m.p. 138-185°C $[\alpha]_D^{25}$ -50.17 (c 0.14, H₂O)
 FAB (C₁₇₄H₂₉₅N₅₅O₅₃): [M+H]⁺ 4005
 AAA: Asx, 2.2(2); Glx, 7.1(7); Ser, 1.7(2); His, 2.8(3); Gly, 1.0(1); Thr, 1.0(1); Ala, 2.1(2); Arg, 3.1(3); Val, 1.1(1); Ile, 1.7(2); Leu, 7.1(7); Lys, 2.7(3).

AVSEHOLLHD KGKSIQDLRR RELLEKLEK LHTRSAW-NH, (SEQ ID NO:72)

Physical Data: m.p. 158-163°C [α]_D²⁵ -46.06 (c 0.17, H₂O)
 FAB (C₁₉₅H₃₂₇N₆₃O₅₅): [M+H]⁺ 4434.8
 AAA: Asx, 2.0(2); Glx, 5.5(6); Ser, 2.7(3); His, 3.1(3); Gly, 1.0(1); Ala,
 1.8(2); Arg, 4.0(4); Thr, 0.9(1); Val, 0.9(1); Ile, 0.9(1); Leu,
 7.5(8); Lys, 3.9(4); Trp, 1.0(1).

-41-

Compound 59:

AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTRSAX-OH (SEQ ID NO:73)
 (X = Nal(2) = 3-(2-naphthyl)-L-alanine)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 10 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr Arg Ser Ala 3-(2-naphthyl)-L-alanine-OH (SEQ ID NO:73)
 35
 15 Physical Data: m.p. 156-162°C $[\alpha]_D^{25}$ -44.44 (c 0.189, H₂O)
 FAB(C₁₉₇H₃₂₈N₆₂O₅₅): [M+H]⁺ 4445.6
 AAA: Asx, 2.1(2); Glx, 5.5(6); Ser, 2.8(3); His, 2.9(3); Gly, 1.0(1); Ala,
 2.0(2); Arg, 4.0(4); Thr, 0.9(1); Val, 1.0(1); Ile, 0.9(1); Leu,
 7.5(8); Lys, 4.2(4); Nal, 1.1(1).

Compound 60:

AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTASAW-OH (SEQ ID NO:74)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 30 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr Ala Ser Ala Trp OH (SEQ ID NO:74)
 35
 35 Physical Data: m.p. 159-164°C $[\alpha]_D^{25}$ -50.94 (c 0.29, H₂O)
 FAB(C₁₉₇H₃₂₀N₆₀O₅₅): [M+H]⁺ 4349.0
 AAA: Asx, 2.0(2); Glx, 5.6(6); Ser, 2.7(3); His, 3.2(3); Gly, 1.0(1); Ala,
 3.1(3); Arg, 2.8(3); Thr, 1.0(1); Val, 1.1(1); Ile, 0.9(1); Leu,
 7.6(8); Lys, 4.0(4); Trp, 1.0(1).

Compound 61:

AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHAEIRA-OH (SEQ ID NO:75)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 50 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr Ala Glu Ile Arg Ala OH (SEQ ID NO:75)
 55 35
 60 Physical Data: m.p. 155-210°C $[\alpha]_D^{25}$ -46.15 (c 0.12, H₂O)
 FAB(C₁₉₅H₃₃₄N₆₀O₅₈): [M+H]⁺ 4475.8
 AAA: Asx, 2.2(2); Glx, 6.9(7); Ser, 1.7(2); His, 3.2(3); Gly, 1.1(1); Ala,
 3.1(3); Arg, 4.0(4); Thr, 0.9(1); Val, 1.1(1); Ile, 1.9(2); Leu,
 8.1(8); Lys, 4.1(4).

-42-

Compound 62:AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTAEIR-OH (SEQ ID NO:76)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 10 Leu His Thr Ala Glu Ile Arg OH (SEQ ID NO:76)
 35

15 Physical Data: m.p. 186-218°C $[\alpha]_D^{25}$ -52.73 (c 0.265, H₂O)
 FAB (C₁₉₂H₃₂₈N₆₁O₅₇): [M+H]⁺ 4404.4
 AAA: Asx, 2.0(2); Glx, 6.6(7); Ser, 1.9(2); His, 3.4(3); Gly, 1.1(1); Ala,
 2.0(2); Arg, 3.8(4); Thr, 1.0(1); Val, 1.1(1); Ile, 1.7(2); Leu,
 7.9(8); Lys, 4.0(4).

20

Compound 63:AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTAEI-OH (SEQ ID NO:77)

25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 30 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr Ala Glu Ile OH (SEQ ID NO:77)
 35

35 Physical Data: m.p. 169-205°C $[\alpha]_D^{25}$ -50.78 (c 0.51, H₂O)
 FAB (C₁₈₆H₃₁₇N₅₇O₅₆): [M+H]⁺ 4248.0
 AAA: Asx, 2.2(2); Glx, 6.8(7); Ser, 1.8(2); His, 3.3(3); Gly, 1.0(1); Ala,
 2.0(2); Arg, 3.0(3); Thr, 1.0(1); Val, 1.0(1); Ile, 1.8(2); Leu,
 7.8(8); Lys, 3.6(4).

40

Compound 64:AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTAE-OH (SEQ ID NO:78)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 50 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr Ala Glu OH (SEQ ID NO:78)
 35

55

Physical Data: m.p. 199-205°C $[\alpha]_D^{25}$ -52.47 (c 0.41, H₂O)
 FAB (C₁₈₀H₃₀₆N₅₆O₅₅): [M+H]⁺ 4135.0
 AAA: Asx, 2.0(2); Glx, 6.6(7); Ser, 1.9(2); His, 3.3(3); Gly, 1.1(1); Ala,
 2.0(2); Arg, 2.9(3); Thr, 1.0(1); Val, 1.1(1); Ile, 1.0(1); Leu,
 8.2(8); Lys, 3.8(4).

60

-43-

Compound 66:SEHQLLHD KGKSIODLRR RELLEKLLEK LHTA-NH₂ (SEQ ID NO:80)

5 Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile Gln Asp
 1 5 10 15
 10 Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys Leu His
 20 25 30
 Thr Ala NH₂ (SEQ ID NO:80)

Physical Data: m.p. 134.2°C $[\alpha]_D^{25}$ -48.12 (c 0.36, H₂O)
 FAB(C₁₆₇H₂₈₆N₅₄O₄₉): [M+H]⁺ 3834.4
 15 AAA: Asx, 2.0(2); Glx, 5.7(6); Ser, 1.7(2); His, 2.9(3); Gly, 1.0(1); Thr,
 0.9(1); Ala, 1.0(1); Arg, 2.8(3); Ile, 0.9(1); Leu, 7.4(8); Lys,
 4.3(4).

20

Compound 67:LLHD KGKSIODLRR RELLEKLLEK LHTA-NH₂ (SEQ ID NO:81)

25 Leu Leu His Asp Lys Gly Lys Ser Ile Gln Asp Leu Arg Arg Arg
 1 5 10 15
 Glu Leu Leu Glu Lys Leu Leu Glu Lys Leu His Thr Ala NH₂ (SEQ ID NO:81)
 20 25

Physical Data: m.p. 128.5-184.5°C $[\alpha]_D^{25}$ -6.53 (c 0.69, MeOH)
 FAB(C₁₄₈H₂₅₉N₄₇O₄₁): [M+H]⁺ 3353
 30 AAA: Asx, 2.0(2); Glx, 4.1(4); Ser, 0.9(1); His, 2.1(2); Gly, 1.0(1); Thr,
 0.9(1); Ala, 1.0(1); Arg, 3.0(3); Ile, 1.0(1); Leu, 8.1(8); Lys,
 35 4.2(4).

Compound 68:LHD KGKSIODLRR RELLEKLLEK LHTA-NH₂ (SEQ ID NO:82)

40 Leu His Asp Lys Gly Lys Ser Ile Gln Asp Leu Arg Arg Arg Glu
 1 5 10 15
 45 Leu Leu Glu Lys Leu Leu Glu Lys Leu His Thr Ala NH₂ (SEQ ID NO:82)
 20 25

Physical Data: m.p. 165-210°C $[\alpha]_D^{25}$ -36.05 (c 0.12, H₂O)
 50 FAB(C₁₄₂H₂₄₈N₄₆O₄₀): [M+H]⁺ 3239.0
 AAA: Asx, 2.0(2); Glx, 3.9(4); Ser, 0.9(1); His, 1.9(2); Gly, 1.0(1); Thr,
 1.0(1); Ala, 1.0(1); Arg, 2.9(3); Ile, 0.9(1); Leu, 6.8(7); Lys,
 4.2(4).

55

- 44 -

Compound 69:SEHQLLHD RGRSIQDLRR RELLERLLER LHAGRRTRSAW-OH (SEQ ID NO:83)

5 Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile Gln Asp
 1 5 10 15
 Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg Leu His
 20 25 30
 10 Leu His Arg Gly Arg Arg Thr Arg Ser Ala Trp OH (SEQ ID NO:83)
 35 40

15 Physical Data: m.p. 150-210°C $[\alpha]_D^{25}$ -18.0 (c 0.64, H₂O)
 FAB (C₂₀₈H₃₅₁N₇₉O₆₀): [M+H]⁺ 4918.6
 AAA: Asx, 2.2(2); Glx, 6.2(6); Ser, 2.8(3); His, 3.1(3); Gly, 2.2(2); Thr,
 2.2(2); Ala, 2.2(2); Arg, 10.4(10); Ile, 1.0(1); Leu, 8.0(8); Trp,
 1.1(1).

20

Compound 70:LLHD RGRSIQDLRR RELLERLLER LHAGRRTRSAW-OH (SEQ ID NO:84)

25 Leu Leu His Asp Arg Gly Arg Ser Ile Gln Asp Leu Arg Arg Arg
 1 5 10 15
 30 Glu Leu Leu Glu Arg Leu Leu Glu Arg Leu His Leu His Arg Gly
 20 25 30
 Arg Arg Thr Arg Ser Ala Trp OH (SEQ ID NO:84)
 35

35 Physical Data: m.p. 150-210°C $[\alpha]_D^{25}$ -41.70 (c 0.36, H₂O)
 FAB (C₁₈₉H₃₂₄N₇₂O₅₂): [M+H]⁺ 4437.14
 AAA: Asx, 2.1(2); Glx, 4.1(4); Ser, 1.9(2); His, 2.0(2); Gly, 2.1(2); Thr,
 2.0(2); Ala, 2.1(2); Arg, 9.7(10); Ile, 0.9(1); Leu, 7.4(8).

40

The side chain cyclized analog (Compound 57) was synthesized as above except the N^ε-Boc-N^ε-Fmoc-Lys and N^ε-Boc-N^ε-Fmoc-Asp were placed in positions 13 and 17, respectively. Upon completion of the Boc amino acid synthesis, the resin was treated with 20% piperidine in DMF at room temperature for 30 minutes. The resin was filtered and washed with DMF, MeOH and CH₂Cl₂. The resin (1.1 g) was suspended in 10 ml DMF containing 250 mg PyBOP. The pH was adjusted to 8-9 with DIEA and the resin stirred for 1 hour. The resin was filtered, washed with DMF and CH₂Cl₂, then resuspended in DMF. The coupling was repeated using 125 mg of PyBOP. The resin was filtered, washed with DMF, MeOH, and CH₂Cl₂, and dried. The resin was then treated with HF and the peptide purified as mentioned above.

-45-

Compound 57:AVSEHQLLHD KGKSIQDLRR RELLEKLLEK LHTA-NH₂ (SEQ ID NO:71)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 10 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr Ala -NH₂ (SEQ ID NO:71)

15 Physical Data: m.p. 142.5-163.5°C [α]_D²⁵ -34.31 (c 0.17, H₂O)
 FAB (C₁₇₅H₂₉₈N₅₆O₅₀): [M+H]⁺ 3986.4
 AAA: Asx, 1.9(2); Glx, 5.9(6); Ser, 1.8(2); His, 3.2(3); Gly, 1.1(1); Ala,
 2.0(2); Arg, 3.0(3); Thr, 1.0(1); Val, 1.1(1); Ile, 0.9(1); Leu,
 8.0(8); Lys, 4.0(4).

20

EXAMPLE II

25 [Met³⁴, Ala³⁵] Compound 1, AVSEHQLLHDKGKSIQDLRRRELLEK-LLEKLHTMA-NH₂,
 (SEQ ID NO:25), was prepared and purified following the procedures of
 Example I. This polypeptide was converted to the homoserine lactone as
 follows. The purified peptide (160 mgs) was dissolved in 44% formic acid
 (4 mL). This solution was combined with a premixed solution of cyanogen
 30 bromide (700 mgs) and phenol (1.6 mgs) in 44% formic acid (4 mL) at 0°C.
 The solution was stirred at 0°C for 2 hr and at room temperature for 2 hrs.
 The formation of the product was monitored by HPLC (Vydac® C-18, 300 Å, 4.6
 x 250 mm, flow of 1.2 mL/min, gradient 25-45% acetonitrile in 0.1% TFA over
 10 min). The reaction was complete within 4 hr. Half of the sample was
 concentrated and purified by preparative RP-HPLC (Vydac® C-18, gradient 25-
 35 45% acetonitrile in 0.1% TFA). The homoserine lactone peptide fractions
 were pooled and lyophilized to give 28 mgs of white solid of >95% purity,
 Compound 4.

Compound 4AVSEHQLLHDKGKSIQDLRRRELLEKLLEKLHTX (X=hSerlac, SEQ ID NO:9)

40 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 45 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr hSerlac (SEQ ID NO:9)

50 Physical Data: m.p. 138-142°C [α]_D²⁵ -50.66° (c 0.1, H₂O)
 FAB (C₁₇₆H₂₉₉N₅₆O₅₂): [M+H]⁺ 4017.61
 AAA: Asp, 2.1(2); Glu, 6.1(6); Ser, 1.8(2); His, 3.0(3); Thi, 1.1(1); Ala,
 1.1(1); Arg, 2.7(3); Val, 1.0(1); Ile, 1.0(1); Leu, 8.2(8); Lys,
 3.8(4); Gly 1.09(1); hSer, 1.09(1).

55

-46-

Similarly, Compound 65 was prepared in accordance with this procedure.

Compound 65:

5 AVSEIOFX₁HN KGKHLSSX₅ER VEWLRKKLOD VHNX₁₅, (SEQ ID NO:79)
 (X₁ = L-norleucine; X₅ = homoserine lactone)

Ala Val Ser Glu Ile Gln Phe Nle His Asn Lys Gly Lys His Leu
 1 5 10 15

Ser Ser Nle Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp
 20 25 30

Val His Asn hSerlac (SEQ ID NO:79)

Physical Data: m.p. 166-176°C [α]_D²⁵ -52.22 (c 0.25, H₂O)

FAB (C₁₈₀H₂₈₈N₅₄O₅₀): [M+H]⁺ 4008.6

AAA: Asx, 3.1(3); Glx, 4.8(5); Ser, 2.9(3); His, 2.9(3); Gly, 1.1(1); Ala, 1.1(1); Arg, 2.0(2); Val, 2.7(3); Phe, 1.0(1); Ile, 1.0(1); Leu + Nle 5.9 (4 + 2); Lys, 2.8(3).

EXAMPLE III

To prepare the homoserine amide, the crude hSerlactone analog, Compound 4, was concentrated and treated with 25 mL saturated NH₃ in methanol. The solution was stirred at 0°C for 2 hr and at room temperature for 16 hr. The reaction was monitored by HPLC (Vydac® C-18, 300 Å, 4.6 x 250 mm, flow of 1.2 mL/min, gradient 20-45% acetonitrile in 0.1% TFA) and was complete within 18 hr. The solution was concentrated and purified by preparative RP-HPLC (Vydac® C-18, gradient of 25-45% acetonitrile in 0.1% TFA). The homoserine amide peptide fractions were pooled and lyophilized to give 30 mgs of white solid of >98% purity, Compound 3.

Compound 3

AVSEHOLLHDKGKSIODLRRRELLEKLLLEKLHTX-NH₂, (X=hSer, SEQ ID NO:8)

40 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15

Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Thr hSer NH₂ (SEQ ID NO:8).

Physical Data:

m.p. 138-142°C

[α]_D²⁵ -45.97° (c 0.25, H₂O)

FAB (C₁₇₆H₃₀₂N₅₆O₅₂): [M+H]⁺ 4033.9

AAA: Asp, 2.1(2); Glu, 6.1(6); Ser, 1.6(2); His, 2.8(3); Gly, 0.97(1); hSer, 0.97(1); Thi, 1.0(1); Ala, 1.0(1); Arg, 2.9(3); Val, 1.0(1); Ile, 1.0(1); Leu, 7.6(8); Lys, 3.9(4).

Similarly, Compounds 22, 23 and 28 were prepared following this procedure.

-47-

Compound 22:

AVSEIOFLHN LGKHLSSLRR RELLEKLEK LHNX-NH₂ (SEQ ID NO:36)
 (X = homoserine)

5 Ala Val Ser Glu Ile Gln Phe Leu His Asn Leu Gly Lys His Leu
 1 5 10 15

10 Ser Ser Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Asn hSer NH₂ (SEQ ID NO:36)

15 Physical Data: m.p. 69.4-128°C $[\alpha]_D^{25}$ -43.93 (c 0.15, H₂O)
 FAB(C₁₇₉H₃₀₂N₅₆O₄₉): [M+H]⁺ 4022.9
 AAA: Asx, 2.0(2); Glx, 4.9(5); Ser, 2.6(3); His, 2.8(3); Gly, 1.0(1); Ala,
 1.0(1); Arg, 3.0(3); Val, 1.0(1); Phe, 1.0(1); Ile, 0.9(1); Leu,
 8.8(9); Lys, 3.4(3);

Compound 23:

AVSEIOFLHN KGKHLSSLRR RELLEKLEK LHNX-NH₂ (SEQ ID NO:37)
 (X = homoserine)

25 Ala Val Ser Glu Ile Gln Phe Leu His Asn Lys Gly Lys His Leu
 1 5 10 15

30 Ser Ser Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Asn hSer NH₂ (SEQ ID NO:37)

35 Physical Data: m.p. 87.1-142.1°C $[\alpha]_D^{25}$ -52.14 (c 0.41, H₂O)
 FAB(C₁₇₉H₃₀₃N₅₇O₄₉): [M+H]⁺ 4038
 AAA: Asx, 2.1(2); Glx, 4.9(5); Ser, 2.7(3); His, 2.8(3); Gly, 1.0(1);
 hSer, 1.0(1); Ala, 1.0(1); Arg, 3.0(3); Val, 1.1(1); Phe, 0.9(1);
 Ile, 0.9(1); Leu, 7.9(8); Lys, 3.7(4).

Compound 28:

AVSEHOLLHD KGKSIODLRR RELLERLLER LHTAGRRX-NH₂ (SEQ ID NO:42)
 (X = homoserine)

45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15

50 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu Glu Arg
 20 25 30

Leu His Thr Ala Gly Arg Arg hSer NH₂ (SEQ ID NO:42)
 35

55 Physical Data: m.p. 80°C $[\alpha]_D^{25}$ -48.64 (c 0.09, H₂O)
 FAB(C₁₉₃H₃₃₄N₇₀O₅₄): [M+H]⁺ 4530.0
 AAA: Asx, 2.2(2); Glx, 6.1(6); Ser, 1.7(2); His, 3.0(3); Gly, 1.9(2);
 hSer, 1.0(1); Thr, 1.0(1); Ala, 2.1(2); Arg, 7.2(7); Val, 0.8(1);
 Ile, 1.0(1); Leu, 8.4(8); Lys, 2.1(2).

60

The homoserine alkylamides were similarly prepared from the homoserin lactone by dissolving it in DMF containing an excess of the corresponding alkylamine. After stirring at room temperature for several

5

AVSEHOLLHD KGKSIQDLRR RELLEKLEK LHTX-NHCH₂CH₃ (SEQ ID NO:69)
(X = homoserine)

20 Physical Data: m.p. (not determined) $[\alpha]_D^{25}$ (not determined)
FAB ($C_{178}H_{306}N_{56}O_{52}$): $[M+H]^+$ 4063.0
AAA: Asx, 2.1(2); Glx, 5.8(6); Ser, 1.7(2); His, 3.1(3); Gly, 0.9(1);
Thr, 1.0(1); Ala, 0.9(1); Arg, 3.0(3); Val, 1.1(1); Ile, 1.0(1); Leu,
8.4(8); Lys, 3.7(4); hSer, 0.9(1).

AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTX-NHCH,CH,C,H. (SEQ ID NO:70)
(X = homoserine)

40 Physical Data: m.p. (not determined) $[\alpha]_D^{25}$ (not determined)
FAB ($C_{184}H_{310}N_{56}O_{52}$): $[M+H]^+$ 4138.8
AAA: Asx, 2.0(2); Glx, 5.9(6); Ser, 1.7(2); His, 2.9(3); Gly, 0.9(1);
Thr, 1.0(1); Ala, 0.9(1); Arg, 3.0(3); Val, 1.0(1); Ile, 0.9(1); Leu,
8.0(8); Lys, 4.1(4); hSer, 0.9(1).

50 The cesium salt of N^ε-Boc-N^α-Fmoc-L-2,4-diaminobutyric acid was attached to Merrifield resin (DMF, 50°C, 48hrs) and used in a solid-phase synthesis in place of the Boc-Ala resin as in Example I. Upon completion of the synthesis the peptide was treated with 20% piperidine in DMF at room temperature for 30 minutes to remove the side chain Fmoc protecting group. The protected peptide spontaneously cyclized to the lactam thereby cleaving 55 itself from the resin. The solution was filtered from the resin and evaporated under vacuum to leave on oil. The residue was treated with liquid HF as in Example I to yield the crude unprotected peptide. The peptide was treated and purified as in Example I.

-49-

Compound 49:

AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTX (SEQ ID NO:63)
(X = L-2,4-diaminobutyryl lactam)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15

10 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Thr L-2,4-diaminobutyryl lactam (SEQ ID NO:63)

15 Physical Data: m.p. 161-181°C $[\alpha]_D^{25}$ -48.38 (c 0.25, H₂O)
 FAB (C₁₇₆H₃₀₀N₅₆O₅₁): [M+H]⁺ 4016.8
 AAA: Asx, 2.1(2); Glx, 6.3(6); Ser, 1.7(2); His, 3.3(3); Gly, 1.1(1); Thr,
 1.0(1); Ala, 2.1(2); Arg, 2.9(3); Val, 0.9(1); Ile, 0.9(1); Leu,
 8.0(8); Lys, 3.8(4).

20

EXAMPLE V

25 An aqueous solution of the homoserine lactone analog from Example II
 was treated with porcine liver esterase (Sigma Chemical Company, St. Louis,
 MO). The hydrolysis of the lactone to the C-terminal homoserine was
 monitored by analytical HPLC. When the hydrolysis was judged to be
 complete the material was purified by preparative HPLC as in Example I.

30

Compound 37:

AVSEHOLLHD KGKSIODLRR RELLEKLLEK LHTX-OH (SEQ ID NO:51)
(X = homoserine)

35 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15

40 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30

Leu His Thr hSer OH (SEQ ID NO:51)

45 Physical Data: m.p. (not determined) $[\alpha]_D^{25}$ (not determined)
 FAB (C₁₇₆H₃₀₁N₅₅O₅₃): [M+H]⁺ 4035.1
 AAA: Asx, 2.1(2); Glx, 5.9(6); Ser, 2.0(2); His, 3.1(3); Gly, 0.8(1);
 hSer, 0.8(1); Thr, 1.0(1); Ala, 1.0(1); Arg, 3.0(3); Val, 1.3(1);
 Ile, 1.0(1); Leu, 8.1(8); Lys, 3.8(4).

50


EXAMPLE VI

55

Following Example I, the protected peptide-resin
 BocAVS(Bzl)E(OBz)H(Bom)QLLHD(OBzl)R(Ts)GR(Ts)S(Bzl)IQD(OBz)-
 LR(Ts)R(Ts)E(OBz)LLE(OBzl)R(Ts)LLK(Fmoc)R(Ts)LH(Bom)T(Bzl)A- O-PAM was
 synthesized on a 0.35 mmol scale. All N^o groups were protected with t-

-50-

butoxycarbonyl (Boc); side chain protecting groups were as indicated. After completion of the synthesis, the peptide resin was treated with 50 mL of 20% piperidine in dimethylformamide (DMF) at room temperature for 30 minutes to remove the fluorenylmethoxy-carbonyl (Fmoc) protecting group on lysine. The resin was washed successively with DMF, MeOH, CH₂Cl₂ and dried to give 1.6 g partially protected peptide. 0.8 g (0.175 mmol) of the partially protected peptide was acylated on lysine with 0.44 g (0.3 mmol) of methoxydi(ethyleneoxy) acetic acid [PEG(2)CH₂COOH] in the presence of 0.16 g (0.3 mmol) benzotriazolyloxy-tris(pyrrolidino)phosphonium hexafluorophosphate (PyBop) and 0.067 g (0.525 mmol) of diisopropylethyl amine (DIEA) in 20 mL DMF at room temperature for 5 hrs. After 5 hrs., the resin was filtered and washed successively with DMF, MeOH and CH₂Cl₂. The acylation step was repeated twice until negative ninhydrin result on the resin was obtained. The final peptide was cleaved from the resin with removal of the side chain protecting groups and purification as in Example I; 100 mgs of Compound 19 were obtained.

Compound 19AVSEHOLLHDRGRSIODLRRRELLERLLKRIHTA-OH (SEQ ID NO:18)

Ala	Val	Ser	Glu	His	Gln	Leu	Leu	His	Asp	Arg	Gly	Arg	Ser	Ile
1				5					10				15	

Gln	Asp	Leu	Arg	Arg	Arg	Glu	Leu	Leu	Glu	Arg	Leu	Leu
			20						25			

Lys(COCH ₂ PEG2)	Arg	Leu	His	Thr	Ala	OH
30						

Physical Data:

m.p. 145-195°C


[α]_D²⁵ -44.60° (c 0.2, H₂O)FAB (C₁₈₃H₃₁₆N₆₄O₅₄): [M+H]⁺ 4276.2

AAA: Asp, 2.1(2); Glu, 5.0(5); Ser, 1.6(2); His, 2.9(3); Gly, 0.9(1); Thr, 1.9(2); Arg, 7.1(7); Val, 1.1(1); Ile, 1.0(1); Leu, 8.0(8); Lys, 0.9(1).

EXAMPLE VII

Peptide was synthesized, cleaved and purified in the same manner as in Example IV except 2-methoxypoly(ethylene-oxy) acetic acid [PEG(5000)CH₂CO₂H] was used as the acylating agent. 0.8 g (0.175 mmol) of partially protected peptide yielded 300 mgs of pure Compound 20.

-51-

Compound 20AVSEHOLLHDRGRSIODLRRRELLERLLKRLHTA-OH (SEQ ID NO:19)

5 Ala Val Ser Glu His Gln Leu Leu His Asp Arg Gly Arg Ser Ile
 1 5 10 15
 10 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Arg Leu Leu
 20 25
 Lys(COCH₂PEG5000) Arg Leu His Thr Ala OH (SEQ ID NO:19).
 30

Physical Data:

15 m.p. 105°C $[\alpha]_D^{25}$ -22.95° (c 0.11, 50% aq. HOAc)
 AAA: Asp, 2.0(2); Glu, 4.8(5); Ser, 1.6(2); His, 2.6(3); Gly, 1.1(1); Thr,
 1.1(1); Arg, 7.3(7); Val, 0.8(1); Ile, 0.9(1); Leu, 8.3(8); Lys,
 1.1(1); Ala, 1.8(2).
 20

EXAMPLE VIIISynthesis of hPTHrp(1-34) analog gene

25 A synthetic gene coding for the hPTHrp(1-34) analog Compound 4 (SEQ
 ID NO:9) was designed having the nucleotide sequence and enzyme restriction
 sites shown in Figure 1. The requisite oligodeoxynucleotides were prepared
 with a DNA synthesizer (Milligen/Bioscience) using the phosphoramidite
 process of Sinha, et al., Nucleic Acid Research 12, 4539-4557 (1984).
 30 After deprotection, the crude oligonucleotides were purified by gel
 electrophoresis on preparative 15% polyacrylamide gels. The
 oligonucleotides were located with UV, excised from the gel, desalted over
 Waters c18 Sep-pak® cartridges, and lyophilized.

35 Amplification via polymerase chain reaction (PCR) was carried out on
 a Perkin-Elmer Cetus thermal cycler, with 25 cycles of: 94°C for 1 minute,
 50°C for 2 minutes, and 72°C for 3 minutes, using reagents, including Taq
 polymerase, from the "GeneAmp" DNA amplification kit (Perkin-Elmer Cetus).

Two overlapping oligonucleotides, an 88mer (2 µg), PTH3,
 (SEQ ID NO:31):

40 CCTCTAGATC TCCGCGGCGC TAGC ATG GCT GTT TCT GAA CAT CAG 45
 Met Ala Val Ser Glu His Gln
 1 5
 CTG CTT CAT GAC AAA GGT AAA TCG ATT CAA GAT CTG AGA CGT C 88
 Leu Leu His Asp Lys Gly Lys Ser Ile Gln Asp Leu Arg Arg
 10 15 20
 45

and an anti-sense 90mer (2 µg), PTH4 (SEQ ID NO:32):

50 CCTCGAAGCT TATGCATCAT TATCTAGA CAT AGT ATG CAG CTT TTC 46
 Met Thr His Leu Lys Glu
 30
 AAG CAG TTT CTC CAG CAG CTC GCG ACG TCT CAG ATC TTG AAT 88
 Leu Leu Lys Glu Leu Leu Glu Arg Arg Leu Asp Gln Ile
 25 20 15
 55 CG 90,

were prepared as the template DNA sequence for the hPTHrp

-52-

(1-34) analog gene. Utilizing the two flanking primers, PTHPCR1: CCTCTAGATC TCCGCGGCGC TAG (SEQ ID NO:33) and PTHPCR2: CCTCGAAGCT TATGCATCAT TATC (SEQ ID NO:34), the entire gene was amplified by PCR. The amplified DNA products were purified by gel electrophoresis on 4% NuSieve® agarose gel. The band containing the synthetic hPTHrp(1-34) analog gene, approximately 150 bases in length, was excised from the gel and approximately 200 ng of DNA isolated by Elu-Quik® glass gel DNA extraction (Schleicher & Schuell, Keene, NH).

EXAMPLE IX

Molecular Cloning of an hPTHrp(1-34)1 Analog Gene

To subclone the hPTHrp(1-34) analog gene of Example VI, 200 ng of the amplified DNA was isolated and cut by restriction enzymes Hind III and Sac II. As shown in Figure 2, the DNA was ligated to 2 µg TrpLE 18 Prot (Ile³,Pro⁵) plasmid previously cleaved with Hind III and Sac II.

The resulting plasmid, TrpLE 18 hPTHrp(1-34)1, containing one copy of the hPTHrp(1-34) analog gene was then transformed into competent *E. coli* HB 101 cells (CLONTECH, Palo Alto, CA). Transformants were subjected to PCR analysis to verify insertion. Transformed cell colonies were selected and boiled in 200 µL of water for 5 minutes; 2 µL were subjected to PCR with two primers flanking the insert. The PCR product was then analyzed on 1% agarose gel to confirm the presence of one copy of the hPTHrp(1-34) gene insert. TrpLE 18 hPTHrp(1-34)1 construct was then verified by DNA sequencing on an automated DNA sequencer (Applied Biosystems Model 373A, Foster City, CA) using the vendor's Dye Deoxy Terminator Sequencing kit.

EXAMPLE X

Construction of a Trp LE 18 vector containing multiple copies of the hPTHrp(1-34) analog gene

Unique Nhe I and Xba I restriction sites are located near the beginning and end of the hPTHrp(1-34) analog gene sequence. These two sites, which recognize different sequences, but produce identical single strand cohesive termini, allow the construction of multiple copy hPTHrp (1-34) genes within the Trp LE 18 vector.

The strategy for constructing repeated hPTHrp(1-34) sequences in tandem is outlined in Figure 3. In separate reactions, 5 µg of plasmid Trp LE 18 hPTHrp(1-34)1 containing a single copy of the gene was cleaved with Bam HI + Nhe I and Xba I + Bam HI. From each digest, about 300 ng of the fragment containing the hPTHrp(1-34) analog gene was isolated. These two fragments were mixed and ligated to form the Trp LE 18 hPTHrp(1-34)2 plasmid. This plasmid was used to transform competent *E. coli* HB 101 cells. Sizing of the transformed PCR products on 1% agarose gel was used to determine the presence of two copies of the hPTHrp(1-34) gene insert. TrpLE 18 hPTHrp(1-34)2 containing two copies of the gene was then confirmed

-53-

by DNA sequencing. The correct fusion of the two hPTHrp(1-34) genes results in the elimination of Nhe I and Xba I sites at the junction. This makes the remaining Xba I and Nhe I sites flanking the tandem genes unique.

By repeating this process the final plasmid Trp LE 18 hPTHrp(1-34)4 containing four copies of the hPTHrp(1-34) gene was constructed, as shown in Figure 4. The sequence of Trp LE 18 hPTHrp(1-34)4 was found to be correct by DNA sequence analysis.

EXAMPLE XI

10 Expression and Purification of Trp LE 18 hPTHrp(1-34)4

Induction of the Trp LE 18 hPTHrp(1-34)4.

A starter culture of 50 mL of LB culture media, J.H. Miller, "Experiments in Molecular Genetics," p.431 (1972), incorporated herein by reference, containing 50µg/mL ampicillin and 100µg/mL tryptophan, was inoculated with *E. coli* cells containing Trp LE 18 hPTHrp(1-34)4 plasmid, and grown overnight at 37°C with vigorous shaking to an A_{550} of about 6. Two liters of LB culture media for production were pre-warmed to 37°C and seeded with 20 mL of the starter culture to give an A_{550} of about 0.06. The culture was then grown with vigorous shaking to an A_{550} of between 0.6 and 0.8, whereupon 2 mL of a 10 mg/mL solution of indole acrylic acid (IAA) was added. Growth was continued with good aeration for about 16 hr to a final A_{550} of about 6 (typically, between 4 and 10). The cells were concentrated by centrifugation and resuspended in 500 mL of 50 mM Tris-HCl, pH 7.5, 0.1 mM EDTA buffer solution (Tris buffer).

25 The suspension was sonicated using a Heat Systems-Ultrasonics, Inc. model 220F sonicator (equipped with a 3/4" horn) operated at 50% of full capacity to avoid overheating.

To determine the extent of induction, the whole cells were analyzed by SDS-PAGE. The gene products derived from the TrpLE 18 hPTHrp(1-34)4 construct were seen as a major band of the predicted MW of approximately 17,000. This accounts for as much as 10% of the total cellular protein.

Isolation of the Fusion Protein.

35 The cell lysate was centrifuged for 15 min. at about 3600 x g to pellet the Trp LE 18 hPTHrp(1-34)4 fusion protein; the supernatant was discarded. The pellet was resuspended in 200 mL Tris buffer. (typically 40-80 A_{550} /mL).

EXAMPLE XII

40 Processing of the Fusion Protein and Purification of homo-Serlactone hPTHrp(1-34) peptide

Cleavage of the methionine residues flanking the hPTHrp(1-34) multimeric fusion protein with CNBr releases the desired homo-Serlactone hPTHrp(1-34) polypeptide, which was purified as described below.

CNBr Treatment of Fusion Protein.

The washed pellet of TrpLE 18 hPTHrp(1-34)₄ fusion protein was resuspended by gently stirring in 60 mL of 70% formic acid (about 20 mg/mL total protein; typically material from 1000 A₅₅₀ units of cells is dissolved in 3 mL). A few drops of octanol were added and N₂ bubbled through the solution for 20 minutes before adding 5.5 g CNBr. This reaction was allowed to proceed for 6 hours at 25°C before an equal volume of 50:50 MeOH:H₂O was mixed with the sample and subsequently removed by rotary evaporation. After 2 to 4 repetitions of this process, the bulk of the formic acid and CNBr were essentially removed. The sample was then evaporated to dryness, redissolved in 200 mL water and lyophilized for storage.

Purification of homo-Serlactone hPTHrp(1-34).

The CNBr cleaved supernatant was dialyzed against 50 mM KH₂PO₄ pH 6.5 for 24 hours with multiple changes. During dialysis, pH was maintained at 6.5. After dialysis, the precipitates were removed by high speed centrifugation. The supernatant was clarified through a Gelman 0.45μ filter device (Acrodisc 4184).

Cation Exchange Chromatography.

Initial purification was accomplished by cation exchange chromatography on a Bio-Gel TSK-SP-5PW HPLC column (21.5 x 150mm). Chromatographic conditions for a flow rate of 8 mL/min and a yield of approximately 12 mg of highly purified homo-Serlactone hPTHrp(1-34) peptide were:

1. Column equilibration in 50 mM KH₂PO₄, pH 6.5
2. Load 10 mL clarified supernatant (approximately 1.5 L culture broth or 2.4 g inclusion).
3. Wash column with 50 mM KH₂PO₄ pH 6.5 containing 50 mM NaCl until baseline is stabilized.
4. Elute column with 50 mM KH₂PO₄ pH 6.5 containing 90 mM NaCl. Collect fractions for about 45 minutes.
5. Analyze the 90 mM NaCl fractions for homo-Serlactone hPTHrp(1-34) content by C18 HPLC before pooling and storage.

Reverse Phase HPLC Chromatography.

A reverse phase Poros R/H 4.6 x 100 mm column (Perseptive Biosystems, Cambridge, MA) was used for the final purification step. The chromatographic conditions were as follows:

-55-

Mobile phase A: 0.1% trifluoroacetic acid (TFA)/water

B: 0.1% trifluoroacetic acid (TFA)/CH₃CN

	<u>TIME</u>	<u>FLOW</u>	<u>% B</u>
5	0 min	4 ml/min	15
	5.0 min	4 ml/min	40
	5.2 min	4 ml/min	100
10	6.8 min	4 ml/min	100
	7.0 min	4 ml/min	15

Retention time of the homo-Serlactone hPTHrp(1-34), Compound 4, was approximately 2.943 minutes. The purified peptide was approximately 98% pure as determined by mass spectroscopy.

EXAMPLE XIII

The compounds of this invention were evaluated for their effect on bone mass in ovariectomized rats, generally in accord with the procedures of Gunness-Hey and Hock, *Metab. Bone Dis.* 5:177 181 (1984).

Adult Sprague-Dawley female rats were acclimatized, weight grouped (n = 9, 10 or 12), and subjected to bilateral ovariectomy (OVX) or sham surgery. Dosing was initiated 17 days after surgery and continued for 20 days. Test compound was administered subcutaneously once a day in 2% rat serum/saline vehicle.

After 20 days of dosing, the rats were sacrificed and the right femurs excised. The femurs were cut in half and the distal half femurs (DHF) were further separated into trabecular bone (TB) and cortical bone (CB) by drilling out the trabeculae. Calcium was extracted and measured by Calcette calcium analyzer and expressed as mean bone Ca in mg/DHF/100 g body weight.

The two sample t-test was used to compare OVX and sham groups. One-way ANOVA was used to compare OVX groups, followed by Fisher's LSD multiple comparison to compare each treatment group to vehicle.

Ovariectomy induced substantial total bone loss, primarily from trabecular bone. Total bone calcium was 47 to 54% lower than for sham-operated controls.

bPTH(1-34) and hPTHrp(1-34) at 80 µg/kg/day provided statistically significant increases in total bone calcium for treated OVX rats, ranging from 53 to 95% and 18 to 40%, respectively; however, there was no significant increase in cortical bone calcium.

Compounds of this invention, dosed at 80 µg/kg/day, increased total bone calcium by from 66 to 138% and trabecular calcium by from 87 to 128%. Cortical bone calcium, trabecular thickness, and bone volume were also significantly increased over untreated OVX controls.

In this assay, the following compounds were tested:

-56-

<u>Compound Number</u>	<u>n (# of tests)</u>	<u>Trabecular Bone Calcium (% increase over OVX)</u>	<u>Total Bone Calcium (% increase over OVX)</u>
Compound 1 (SEQ ID NO:7)	6	101-128%	88-138%
Compound 2 (SEQ ID NO:6)	3	87-102%	66-114%
Compound 4 (SEQ ID NO:9)	3	-	88-114%

10 In similar studies, ovariectomized rats were dosed for 5, 10 and 20 days, at 40 ug/kg/day, with the following results:

<u>Compound Number</u>	<u>n (# of tests)</u>	<u># of Days (d) of dosing</u>	<u>Total Bone Calcium (% increase over OVX)</u>
Compound 1 (SEQ ID NO:7)	3	20d	73-109%
Compound 4 (SEQ ID NO:9)	5	20d	79-105%
Compound 4 (SEQ ID NO:9)	1	10d	79%
Compound 49 (SEQ ID NO:63)	1	10d	93%
Compound 4 (SEQ ID NO:9)	1	5d	55%
Compound 42 (SEQ ID NO:56)	1	5d	60%

25

EXAMPLE XIV TOXICITY

In the above Example XI, no toxic effects were observed with the compounds of the invention.

-57-

We claim:

1. A synthetic polypeptide analog of parathyroid hormon (PTH), parathyroid hormone related peptide (PTHrp), or of a physiologically active truncated homolog or analog of PTH or PTHrp, or salt thereof, in which amino acid residues (22-31) form an amphipathic α -helix, said residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa), ordered in the sequence:

Haa (Laa Laa Haa Haa)₂ Laa.

2. A polypeptide of claim 1, or salt thereof, in which amino acid residues (22-31) form an amphipathic α -helix, the sequence of said residues (22-31) selected from:

a) Xaa¹ Xaa² Leu Xaa⁴ Xaa⁵ Leu Xaa⁷ Xaa⁸ Xaa⁹ Xaa¹⁰ wherein

Xaa¹ and Xaa⁴ are independently Glu, Glu(OCH₃), His, or Phe; Xaa² is Leu or Phe; Xaa⁵ is Lys or His; Xaa⁷ and Xaa¹⁰ are independently Leu or Ile; Xaa⁸ is Ala, Arg, or Glu; and Xaa⁹ is Lys or Glu (SEQ ID NO:85); preferably

Glu Leu Leu Glu Lys Leu Leu Xaa Lys Leu wherein

Xaa is Glu or Arg (SEQ ID NO:26);

b) Xaa¹ Xaa² Leu Xaa⁴ Arg Leu Leu Xaa⁸ Arg Leu wherein

Xaa¹ and Xaa⁴ are independently Glu, Glu(OCH₃), His, or Phe; Xaa² is Leu or Phe; Xaa⁸ is Glu, Lys, or Lys(COCH₂PEGX) and PEGX is a poly-(ethylene glycol methyl ether) radical of molecular weight 100 to 10,000 (SEQ ID NO:86); preferably,

Glu Leu Leu Glu Arg Leu Leu Xaa Arg Leu wherein

Xaa is Glu, Lys, or Lys(COCH₂PEGX) and PEGX is a poly-(ethylene glycol methyl ether) radical of molecular weight 100 to 10,000 (SEQ ID NO:27);

c) Ala Leu Ala Glu Ala Leu Ala Glu Ala Leu (SEQ ID NO:28);

d) Ser Leu Leu Ser Ser Leu Leu Ser Ser Leu (SEQ ID NO:29);

e) Ala Phe Tyr Asp Lys Val Ala Glu Lys Leu (SEQ ID NO:30).

3. A polypeptide of claim 2 of the formula:

Xaa¹ Xaa² Xaa³ Xaa⁴ Xaa⁵ Xaa⁶ Xaa⁷ Leu His Xaa¹⁰ Xaa¹¹ Gly Xaa¹³ Ser Ile Gln Xaa¹⁷ Leu Xaa¹⁹ Xaa²⁰ Xaa²¹ Xaa²²⁻³¹ Xaa³² Xaa³³ Xaa³⁴ Xaa³⁵ Xaa³⁶ Xaa³⁷ Xaa³⁸ Term, wherein:

Xaa¹ is absent or is Ala;

Xaa² is absent or is Val;

Xaa³ is absent or is Ser;

Xaa⁴ is absent or is Glu or Glu(OCH₃);

Xaa⁵ is absent or is His or Ala;

-58-

Xaa⁶ is absent or is Gln;
 Xaa⁷ is absent or is Leu;
 Xaa¹⁰ and Xaa¹⁷ are independently Asp or Asp(OCH₃);
 Xaa¹¹ is Lys, Arg, or Leu;
 5 Xaa¹³ is Lys, Arg, Tyr, Cys, Leu, Cys(CH₂CONH(CH₂)₂NH(biotinyl)), Lys(7-dimethylamino-2-oxo-2H-1-benzopyran-4-acetyl), or Lys(dihydrocinnamoyl);
 Xaa²⁰ is Arg or Leu;
 Xaa¹⁹ and Xaa²¹ are independently Lys, Ala, or Arg;
 Xaa²²⁻³¹ is selected from (SEQ ID NOS:26, 27, 28, 29, or 30); Xaa³² is His,
 10 Pro, or Lys;
 Xaa³³ is absent, or is Pro, Thr, Glu, or Ala;
 Xaa³⁴ is absent, or is Pro, Arg, Met, Ala, hSer, hSer lactone, Tyr, Leu, or 1,4-diaminobutyryl lactam;
 Xaa³⁵ is absent or is Pro, Glu, Ser, Ala, or Gly;
 15 Xaa³⁶ is absent or is Ala, Arg, or Ile;
 Xaa³⁷ is absent or is Arg, Trp, or 3-(-2-naphthyl)-L-alanine;
 Xaa³⁸ is absent or is Ala or hSer or Xaa³⁸⁻⁴² is Thr Arg Ser Ala Trp;
 and Term is OR or NR₂ where each R is independently H, (C₁-C₄)alkyl or phenyl(C₁-C₄)alkyl; and the pharmaceutically acceptable salts thereof.

20
 4. A polypeptide of claim 3 wherein Xaa²²⁻³¹ is (SEQ ID NO:26), and wherein Xaa¹¹ and Xaa¹³ are both Lys; and Xaa¹⁹ and Xaa²¹ are both Arg.

5. The polypeptide of claim 4 which is:
 25 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 30 20 25 30
 Leu His Thr Ala NH₂ (SEQ ID NO:7).

6. The polypeptide of claim 4 which is:
 35 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 40 20 25 30
 Leu Pro NH₂ (SEQ ID NO:56).

7. The polypeptide of claim 4 which is:
 45 Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 50 20 25 30
 Leu His Thr 1,4-diaminobutyryl lactam (SEQ ID NO:63).

-59-

8. The polypeptide of claim 4 which is:

Ala Val Ser Glu His Gln Leu Leu His Asp Lys Gly Lys Ser Ile
 1 5 10 15
 Gln Asp Leu Arg Arg Arg Glu Leu Leu Glu Lys Leu Leu Glu Lys
 20 25 30
 Leu His Thr hSerlac (SEQ ID NO:9).

9. A pharmaceutical composition comprising an effective bone mass increasing amount of a polypeptide of claim 1, or salt thereof, in admixture with a pharmaceutically acceptable carrier.

10. A method for treating mammalian conditions characterized by decreases in bone mass, which method comprises administering to a subject in need thereof an effective bone mass increasing amount of a synthetic polypeptide analog of parathyroid hormone (PTH), parathyroid hormone related peptide (PTHrp), or of a physiologically active truncated homolog or analog of PTH or PTHrp, or salt thereof, in which amino acid residues (22-31) form an amphipathic α -helix, said residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa), ordered in the sequence:

Haa (Laa Laa Haa Haa)₂ Laa.

11. A method of claim 10 in which the condition to be treated is osteoporosis.

12. A process for the solid phase synthesis of a polypeptide analog of PTH, PTHrp, or of a physiologically active truncated homolog or analog of PTH or PTHrp, or salt thereof, in which amino acid residues (22-31) form an amphipathic α -helix, said residues (22-31) selected from hydrophilic amino acids (Haa) and lipophilic amino acids (Laa), ordered in the sequence:

~~Haa-(Laa-Laa-Haa-Haa)₂-Laa,~~

which process comprises sequentially coupling protected amino acids on a suitable resin support, removing the side chain and N^o protecting groups, and cleaving the polypeptide from the resin support.

13. The use of a compound of Claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of mammalian conditions characterized by decreases in bone mass, particularly, for the treatment of osteoporosis.






Figure 2

SUBSTITUTE SHEET
ISA/EP




Figure 3

SUBSTITUTE SHEET
ISA/EP




Figure 4

SUBSTITUTE SHEET
ISA/EP

International Application No

International Application No. 93/08463		
I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶ According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 C07K7/10; A61K37/43		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07K ; A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ^o	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	EP,A,0 451 867 (MITSUBISHI KASEI CORPORATION) 16 October 1991 see the whole document ---	1,2,9-13
X	EP,A,0 477 885 (TAKEDA CHEMICAL INDUSTRIES,LTD) 1 April 1992 see page 5, line 54; claims 1-7,9; example 21 ---	1,9-13
P,X	WO,A,9 306 845 (P.PANG ET AL) 15 April 1993 SEQ ID No 6,8,10; CLAIMS 1-10; FIG. 3,5,7,9 ---	1,9-13
	-/--	
^o Special categories of cited documents : ¹⁰ "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
22 OCTOBER 1993	23 -11- 1993	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	GROENENDIJK M.S.	

Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	<p>BIOCHEMISTRY. vol. 31, no. 7, 25 February 1992, EASTON, PA US pages 2056 - 2063 W.NEUGEBAUER ET AL 'STRUCTURAL ELEMENTS OF hPTH AND THEIR POSSIBLE RELATION TO BIOLOGICAL ACTIVITIES' see the whole document</p> <p style="text-align: center;">---</p>	1-13
A	<p>US,A,4 086 196 (G.F.TREGAR) 25 April 1978 see the whole document</p> <p style="text-align: center;">-----</p>	1-19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/06465

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark : Although claims 10 and 11 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/ composition.
2. ☒ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The definitions of "hydrophilic" and "lipophilic" in claim 1 seem to be rather arbitrary and inconsequent, e.g. Ala is considered to belong to both categories and she is apparently considered to be hydrophilic (see cl.2,p. 57,1.16). Consequently the search has been restricted to the scope of cl.2.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9306465
SA 76687

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 22/10/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0451867	16-10-91	JP-A- 4217997 US-A- 5229489	07-08-92 20-07-93
EP-A-0477885	01-04-92	JP-A- 5032696	09-02-93
WO-A-9306845	15-04-93	AU-A- 2871292	03-05-93
US-A-4086196	25-04-78	None	

EPO FORM P0079

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)